

Longitudinal Effects of Prenatal Alcohol Exposure on Visual Neurodevelopment Over Infancy

Emma T. Margolis¹, Lauren Davel², Niall J. Bourke³, Cara Bosco¹, Michal R. Zieff², Alexa D. Monachino¹, Thandeka Mazubane², Simone R. Williams², Marlie Miles², Chloë A. Jacobs², Sadeeka Williams², Layla Bradford², Candice Knipe², Zamazimba Madi², Bokang Methola², Tembeka Mhlakwaphalwa², Nwabisa Mlandu², Khanyisa Nkubungu², Zayaan Goolam Nabi², Tracy Pan², Reese Samuels², Nicolò Pini⁴, Vanja Klepac-Ceraj⁵, William P. Fifer⁴, Daniel C. Alexander⁶, Derek K. Jones⁷, Steve C. R. Williams³, Dima Amso⁸, Kirsten A. Donald², and Laurel J. Gabard-Durnam¹

¹ Department of Psychology, Northeastern University

² Department of Paediatrics and Child Health, University of Cape Town

³ Department of Neuroimaging, King's College London

⁴ New York State Psychiatric Institute, Irving Medical Center, Columbia University

⁵ Department of Biological Sciences, Wellesley College

⁶ Department of Computer Science, University College London

⁷ School of Psychology, Cardiff University

⁸ Department of Psychology, Columbia University

This article was published Online First March 21, 2024.

Emma T. Margolis  <https://orcid.org/0000-0002-2036-8078>

This research is supported by funding from the Wellcome Leap 1kD Program to Vanja Klepac-Ceraj, William P. Fifer, Daniel C. Alexander, Derek K. Jones, Steve C. R. Williams, Dima Amso, Kirsten A. Donald, and Laurel J. Gabard-Durnam. The views expressed in this article are those of the authors and do not necessarily represent those of the Wellcome Leap 1kD Program. The authors would like to thank the mothers, families, and infants who made this work possible. The authors would also like to thank the Khula recruitment team in South Africa (especially site coordinator, Ringie Gulwa, and fieldworker, Pamela Madikane) for their work in recruiting and connecting with the families in this study. This study was not preregistered. Scripts for the primary and exploratory analyses reported are available from the Open Science Framework (https://osf.io/kjtsu/?view_only=30b08cd27a0f412cbff10dddf39561fd). Data may be made available upon reasonable request.

Kirsten A. Donald and Laurel J. Gabard-Durnam contributed equally to this work and share the last authorship. Emma T. Margolis served as lead for conceptualization, formal analysis, methodology, visualization, writing—original draft, and writing—review and editing and served in a supporting role for data curation. Lauren Davel served as lead for data curation and investigation and served in a supporting role for writing—review and editing. Niall J. Bourke served as lead for data curation, contributed equally to formal analysis, writing—original draft, and served in a supporting role for writing—review and editing. Cara Bosco served as lead for data curation and project administration and served in a supporting role for writing—review and editing. Michal R. Zieff served as lead for data curation, investigation, and project administration and served in a supporting role for writing—review and editing. Alexa D. Monachino served as lead for software and served in a supporting role

for writing—review and editing. Thandeka Mazubane served as lead for data curation and investigation and served in a supporting role for writing—review and editing. Simone R. Williams served as lead for data curation and investigation and served in a supporting role for writing—review and editing. Marlie Miles contributed equally to investigation and project administration and served in a supporting role for writing—review and editing. Chloë A. Jacobs, Sadeeka Williams, Layla Bradford, Candice Knipe, Zamazimba Madi, Bokang Methola, Tembeka Mhlakwaphalwa, Nwabisa Mlandu, Khanyisa Nkubungu, Zayaan Goolam Nabi, Tracy Pan, and Reese Samuels contributed equally to investigation and served in a supporting role for writing—review and editing. Nicolò Pini, Vanja Klepac-Ceraj, William P. Fifer, Daniel C. Alexander, Derek K. Jones, Steve C. R. Williams, and Dima Amso served in a supporting role for writing—review and editing. Kirsten A. Donald served as lead for supervision and contributed equally to writing—review and editing. Laurel J. Gabard-Durnam served as lead for conceptualization, formal analysis, methodology, software, writing—original draft, and writing—review and editing. Vanja Klepac-Ceraj, William P. Fifer, Daniel C. Alexander, Derek K. Jones, Steve C. R. Williams, Dima Amso, and Kirsten A. Donald contributed equally to methodology. Vanja Klepac-Ceraj, William P. Fifer, Daniel C. Alexander, Derek K. Jones, Steve C. R. Williams, Dima Amso, Kirsten A. Donald, and Laurel J. Gabard-Durnam contributed equally to funding acquisition.

Correspondence concerning this article should be addressed to Emma T. Margolis, Department of Psychology, Northeastern University, 805 Columbus Avenue, Boston, MA 02120, United States. Email: margolis.emm@northeastern.edu

Prenatal alcohol exposure (PAE) affects neurodevelopment in over 59 million individuals globally. Prior studies using dichotomous categorization of alcohol use and comorbid substance exposures provide limited knowledge of how prenatal alcohol specifically impacts early human neurodevelopment. In this longitudinal cohort study from Cape Town, South Africa, PAE is measured continuously—characterizing timing, dose, and drinking patterns (i.e., binge drinking). High-density electroencephalography (EEG) during a visual-evoked potential (VEP) task was collected from infants aged 8 to 52 weeks with prenatal exposure exclusively to alcohol and matched on sociodemographic factors to infants with no substance exposure in utero. First trimester alcohol exposure related to altered timing of the P1 VEP component over the first 6 months postnatally, and first trimester binge drinking exposure altered timing of the P1 VEP components such that increased exposure was associated with longer VEP latencies while increasing age was related to shorter VEP latencies ($n = 108$). These results suggest alcohol exposure in the first trimester may alter visual neurodevelopmental timing in early infancy. Exploratory individual-difference analysis across infants with and without PAE tested the relation between VEP latencies and myelination for a subsample of infants with usable magnetic resonance imaging (MRI) T1w and T2w scans collected at the same time point as EEG ($n = 47$). Decreased MRI T1w/T2w ratios (an indicator of myelin) in the primary visual cortex ($n = 47$) were linked to longer P1 VEP latencies. Results from these two sets of analyses suggest that prenatal alcohol and postnatal myelination may both separately impact VEP latency over infancy.

Public Significance Statement

For infants prenatally exposed exclusively to alcohol, continuous measures of first trimester exposure and binge drinking exposure were associated with altered timing of visual neurodevelopment over the first 6 months postnatally. Myelination may serve as a partial mechanism for these differences. The World Health Organization Alcohol, Smoking, and Substance Involvement Screening Test is an acceptable tool to detect alcohol risk in pregnant populations. This work can inform efforts to support pregnant people and their infants pre- and postnatally.

Keywords: prenatal alcohol exposure, infant neurodevelopment, fetal programming, visual-evoked potential event-related potential, myelination

Supplemental materials: <https://doi.org/10.1037/dev0001727.supp>

Alcohol is a known teratogen that affects central nervous system development in utero (Coles, 1994; Jones & Smith, 1973; Lemoine et al., 1968). Globally, 9.8% of pregnant people endorse alcohol use during their pregnancy, affecting neurodevelopment in over 59 million individuals (Popova et al., 2018). Rates and patterns of prenatal alcohol exposure (PAE) vary by country, culture, and across communities. Certain communities report high incidences of PAE, including in South Africa (Popova et al., 2023). A recent study of 5,231 women receiving prenatal care and delivery in the Cape Town metropolitan area, South Africa found that up to 36.9% of the sample self-reported alcohol use during pregnancy and/or in the 3 months before pregnancy recognition (Petersen Williams et al., 2014).

Contextualizing PAE in South African Communities

It is important to recognize the structural and cultural factors, including historical sociopolitical frameworks, which might contribute to the high prevalence of PAE present in certain South African communities. The colonial apartheid regime in South Africa created a system of significant economic and sociopolitical inequalities that served to disadvantage communities such as those in the current study sample (Adebiyi et al., 2021; Jacobson et al., 2017; Marshall et al., 2022; May et al., 2019). Additionally, in health systems with access to limited resources, more acutely life-threatening medical conditions may be prioritized over conditions that are more chronic and less immediately life-threatening, leading to less resource allocation to mental health and substance use disorder treatment and support

(Williams & Wyatt, 2015). From an intersectional perspective (Crenshaw, 1989), some work suggests that intersecting identities for Black women in South Africa may lead to additional internal, external, and structural barriers in accessing, receiving, and completing treatment for alcohol use disorders (Pretorius et al., 2009). On an individual level, multiple explanations for alcohol use during pregnancy emerged from qualitative interviews conducted with peripartum women in informal settlements in the Cape Town metropole (Fletcher et al., 2018; Watt et al., 2014). Notably, one of the major themes identified was using drinking as a way to cope with stressors, abuse, or other trauma, both those acutely related to the pregnancy and new child or other unrelated stressors. Women in both studies also cited drinking as a way to socialize, drinking during pregnancy being an acceptable social norm in their communities, and lack of attachment to the pregnancy or not wanting the pregnancy as reasons for their drinking.

PAE and Differences in Visual Processing During Childhood

PAE may result in a spectrum of physical, sensory, and/or learning disabilities for exposed individuals. The spectrum of disabilities is commonly referred to as fetal alcohol spectrum disorders (FASD). Behavioral literature in school-aged children has established sensory differences as consistent outcomes of prenatal exposure, notably with respect to visual processing (National Institute on Alcohol Abuse and Alcoholism, 2000). Preschool and early school-age children

with PAE have observed difficulties in visual-motor integration skills (Janzen, 1995; Mattson et al., 1998). Additional studies observed difficulties in visual-spatial processing and functioning in 9- to 16-year-old children with PAE (Hunt et al., 1995; Mattson et al., 1996; Uecker & Nadel, 1996). One behavioral study conducted in infancy found alterations in visual fixation behavior for infants with PAE, which might be indicative of these visual differences emerging early in development (Jacobson et al., 1993). However, limited studies in human development examine brain phenotypes that may link the prenatal exposure to these later outcomes (Coles & Li, 2011; Donald et al., 2015; Wozniak & Muetzel, 2011).

Not only is early visual processing more obviously indicative of later visual-related abilities (Harter et al., 1977; Hartmann, 1995; McCulloch, 2013; Taylor et al., 1992), additional evidence suggests that early sensory processing, most notably in the visual domain, scaffolds later executive functioning via visual attention pathways (Amso & Scerif, 2015; Rosen et al., 2019; Werchan & Amso, 2017). Therefore, it is imperative to understand the emerging visual system in infants with PAE to better support later visual and cognitive development.

Impact of PAE on Visual Neuroplasticity Mechanisms

One potential mechanism through which PAE may impact postnatal sensory development is via neuroplasticity regulation for sensory learning over the first year of life. Increased neuroplasticity results in rapid sensory learning during this window, which can be assessed in the human both functionally and structurally using noninvasive, in vivo techniques such as electroencephalography (EEG; Barry-Anwar et al., 2020) and magnetic resonance imaging (MRI; Deoni et al., 2011; Goddings et al., 2021), respectively (Gabard-Durnam & McLaughlin, 2020).

In humans, this visual learning and plasticity can be indexed neurally through a visual-evoked potential (VEP) task using EEG (Barry-Anwar et al., 2020). This type of visual Event-Related Potential (ERP) provides a time-locked response to a visual stimulus which can be used as a marker of the underlying learning and development (Lippé et al., 2006, 2009). For example, as learning accumulates and the brain develops, sensory processing speeds up and neural responses come to resemble mature spatiotemporal patterns (Barry-Anwar et al., 2020). Specifically, the VEP comprises three main components: a negative N1 peak, a positive P1 peak, and a negative N2 peak. As infants gain more visual experience and their brain develops, these VEP components occur more quickly in response to visual stimuli; that is, the latency to each component's peak amplitude becomes shorter (Lippé et al., 2009). These changes occur rapidly over infancy, with the adult pattern of response emerging around 24 months of age (Lippé et al., 2009). The VEP response (when measured via magnetoencephalography) as used in this study has been localized to activity within the visual cortex from primarily two sources, the primary visual cortex (V1; components N1, P1, N2) and the middle temporal visual area (i.e., MT or V5; components P1, N2; Barnikol et al., 2006).

Prior studies in human infancy suggest the VEP in early life is sensitive to PAE. A study with a sample of human infants with heterogeneous prenatal substance exposures found that infants with more first trimester alcohol exposure (quantified as drinks per week within the first trimester) demonstrated prolonged P1 VEP latencies at 1 month of age when compared to those without first trimester alcohol exposure (Scher et al., 1998). Analyses examining second or third trimester use in that same study did not find the same effects on the P1 latency as the

first trimester exposure (though other substances also impacted the P1 latency), indicating the need to examine prenatal alcohol use in terms of both timing and dose with respect to VEP development. Although the heterogeneous substance use precludes conclusions about the specificity of this effect to alcohol, and effects of subsequent trimesters may have been confounded by prior trimester(s) exposure (unaccounted for in models), the longer P1 latencies observed are consistent with delayed early visual development following alcohol exposure. Another study in neonates with PAE has also demonstrated differential electrophysiological responses following a visual stimulus when compared to neonates without exposure, though exposure was quantified categorically rather than by dose (Olegård et al., 1979). Indeed, the majority of prior literature on PAE has measured exposure categorically (comparing those with high exposure levels to those with no prenatal exposure (Coles & Li, 2011) so dose-related effects as a function of trimester timing remain poorly understood despite early evidence this approach matters for interrogating visual neurodevelopment.

Visual plasticity is also regulated by structural changes in myelin over early development that can be detected with MRI. Specifically, myelination happens most rapidly in infancy, with occipital lobes largely myelinated by 5–6 months (Dean et al., 2014). Among its many other roles, myelination serves as an active player in the regulation of sensory neuroplasticity by acting as both a “brake” on this plasticity and a “shield” protecting prior functional learning from future insults (e.g., by preserving the structural connections serving learned circuit function; Nelson & Gabard-Durnam, 2020). Myelination also increases nerve conduction velocity and is therefore associated with changes in VEP latency in clinical populations and in animal models at other life stages, such that increased myelination is associated with shorter latencies (Heidari et al., 2019; Klistorner & Graham, 2021; Walt et al., 2015). It has long been hypothesized that increased myelination in the visual cortex over infancy is also partly responsible for the latency shifts in VEP over infancy, thus indicating preservation of the visual learning and efficient visual cortex responsiveness.

Animal model literature and human literature beginning in childhood have characterized the effects of alcohol exposure on myelination (e.g., Almeida et al., 2020; Darbinian & Selzer, 2022; Milbocker et al., 2022). Human histological studies also provide evidence for the impact of PAE on dysmyelination (Darbinian & Selzer, 2022). Human neuroimaging studies using MRI found white matter microstructural alterations, including in the occipital lobe, in individuals as young as 5 years of age with PAE and/or FASD (Donald et al., 2015; Ghazi Sherbaf et al., 2019; McLachlan et al., 2019; Wozniak & Muetzel, 2011). However, it is important to characterize these microstructural differences in infancy, during critical windows of myelination and functional learning to understand how and when these differences emerge.

In summary, sensitive periods are tightly regulated by biological mechanisms including by “brakes” which punctuate these sensitive periods and preserve/protect the learning that has occurred (Gabard-Durnam & McLaughlin, 2020; Hensch & Bilimoria, 2012; Reh et al., 2020; Takesian & Hensch, 2013). Exposure to alcohol prenatally impacts sensitive period regulators such as myelination (Almeida et al., 2020; Darbinian & Selzer, 2022; Donald et al., 2015; Wozniak & Muetzel, 2011), which is a key “brake” on sensitive periods. By disrupting regulators such as myelin, alcohol exposure may shift the timing of postnatal sensitive periods, leading to downstream, potentially disordered consequences on cognition

and behavior (e.g., Hensch, 2005; Ivanov, 2021; Makinodan et al., 2012; Marín, 2016; Takesian & Hensch, 2013).

The Importance of Considering Timing and Dose Effects of PAE

It is important to consider the substantial heterogeneity in presentation for individuals with PAE. Timing, dose, and drinking pattern differences contribute to this heterogeneity in postnatal disabilities and underlying neural substrates (O'Leary et al., 2010; Pini et al., 2019; Subramoney et al., 2018). Trimester timing effects are perhaps the best characterized of these factors to date, in which those with PAE during first trimester organogenesis display craniofacial abnormalities such as thin upper lip, smooth philtrum, and short palpebral fissure (Popova et al., 2023), and those with exposure after 8 weeks gestation are less likely to present with this facial phenotype that is typically associated with PAE. However, those with later exposure are not immune to the effects of alcohol and exhibit alterations in brain development. Notably, delayed P1 VEP latencies at 1 month of age have been found in those with alcohol exposure in the first trimester (Scher et al., 1998).

Multiple studies of human behavior have characterized dose-related effects of PAE (e.g., Smith et al., 1986; Streissguth et al., 1989; Testa et al., 2003), but few studies examining neurodevelopment in humans have examined PAE with dose-specific information (Almeida et al., 2020; Jacobson et al., 2017). One neurodevelopmental study in 10-year-old children with heterogeneous prenatal substance exposure found that increased PAE dose was linked to greater cerebral volume and had nonlinear associations with regional brain volumes including the occipital lobes (Lees et al., 2020). In a more general sense, the majority of human studies use clinical samples and therefore highlight the effects of PAE at high doses, but there is a need for more human work characterizing the effects at more moderate levels of PAE (Dejong et al., 2019).

There is also evidence that exposure matters not only in terms of timing or amount, but also in the way in which these two factors interact to produce different drinking patterns. Binge drinking for women, according to the U.S. Centers for Disease Control and Prevention, is defined as drinking four or more drinks in a drinking episode (CDC, 2022). Animal models observed more severe consequences with regards to brain growth (i.e., more severe microcephaly) for rats with alcohol exposure following a binge drinking pattern as opposed to those exposed to an equal amount of alcohol following a more uniform pattern (Kelly et al., 1987; West et al., 1987). Other studies have reported increased difficulties in regulation of executive functioning for school-age children with prenatal binge drinking exposures (Bailey et al., 2004; Jacobson et al., 2021; Nulman et al., 2004). However, it remains unknown how binge exposures impact human neurodevelopment underlying these behavioral differences. Thus, in studies investigating the effects of PAE, it is imperative to consider timing across trimesters, dose effects, and pattern of drinking exposure (i.e., binge drinking), as these may differentially impact the neural substrates of early development.

Current Study

The current study addresses limitations in the current literature regarding how and when PAE is biologically embedded to shape

postnatal brain development and result in sensory behavioral outcomes using longitudinal data from an ongoing project with 394 families recruited from Gugulethu, an informal settlement, in Cape Town, South Africa. This study may contribute to the field's understanding of how alcohol teratogenesis impacts postnatal neurodevelopment in several ways. First, PAE was quantified by trimester (to account for fluctuations across trimesters) using a continuous rather than dichotomous index. Distinguishing alcohol exposures by trimester facilitates the investigation of how the timing, dose, and pattern of PAE engender teratogenesis. Second, the homogeneous alcohol exposure measured in the current sample removes confounding effects of substance comorbidity found in prior studies. Furthermore, the multimodal longitudinal design of the current study allows for indexing alcohol's effects on functional neurodevelopmental changes via EEG VEP components, and exploring potential structural changes (via MRI T1w/T2w ratios to estimate myelination; Glasser & Van Essen, 2011) that may in part contribute to functional neurodevelopment during the period of rapid development that has been historically understudied. Specifically, this study set out to test (a) how PAE (i.e., timing, dose, and binge behavior) impacts electrophysiological responses to visual stimuli, as indexed by latencies of the EEG VEP components. We predicted that earlier exposure as well as increased dose and binge episodes would result in longer VEP latencies, indicating delayed neurodevelopment or less-efficient visual learning. We also investigated (b) how changes in VEP component latencies in this sample relate to underlying differences in myelination during this critical window. We predicted that shorter latencies would be associated with increased myelination of the primary visual cortex (V1) and middle temporal visual area (MT/V5), as indexed by T1w/T2w ratios from MRI.

Method

Participants

Participants for the current study come from an ongoing prospective longitudinal project with 394 mothers and their infants recruited from Gugulethu, an informal settlement, in Cape Town, South Africa (for study description, see Zieff et al., under review). The first language of the majority of residents in this area is Xhosa. Three hundred twenty-nine mothers were recruited prenatally, and 65 mothers were recruited postnatally from local community clinics. One infant was excluded after birth using a priori exclusion criteria due to extreme prematurity (born 8 weeks premature). This study was approved by the relevant university Health Research Ethics Committees. Informed consent was collected from mothers on behalf of themselves and their infants.

Study Design

All questionnaires were administered by an interviewer (trained study nurse or research assistant). Questionnaires and procedures were offered/explained in English or Xhosa depending on language preference of the mother. At the enrollment study visit, mothers completed a series of questionnaires including the Alcohol Exposure Questionnaire (AEQ; Kwiatkowski et al., 2018), World Health Organization Alcohol, Smoking, and Substance Involvement Screening Test (WHO ASSIST; WHO ASSIST Working Group, 2002), and Edinburgh Postnatal Depression Scale (EPDS; Cox et al., 1987). Demographic information including maternal place of birth,

primary spoken language, maternal age at enrollment, maternal educational attainment, and maternal income were also collected at the enrollment visit.

Families were invited to participate in two in-lab study visits over their infant's first year of life, one visit in the first 6 months and another visit from 6 to 12 months of age. At the first in-lab study visit (hereafter Visit 1), occurring when infants were between approximately 2 and 6 months of age (age in weeks: $M = 15.798$, $SD = 3.719$, range = 8.286–24.571), the following data were collected: the infants' age (in weeks), sex, infant EEG, and MRI. For infants who were unable to complete both EEG and MRI on the same day, EEG, and MRI were collected on different days (for those with multimodal data included in the current study, median age difference = 1.571 weeks).

At the second study visit (hereafter Visit 2), occurring when infants were between approximately 6 months and 12 months of age (age in weeks: $M = 36.489$, $SD = 6.337$, range = 26.000–51.857), infant EEG and MRI data were collected again. Not all infants had EEG and MRI data collected at both timepoints or contributed usable data at both timepoints.

Prior work has explored validity of the standardized clinical cutoffs of the EPDS (Cox et al., 1987) within the context of diverse African and South African communities (Donald et al., 2019; Lawrie et al., 1998; Pellowski et al., 2023; Tsai et al., 2013) and has validated the scale in the Xhosa language (Lawrie et al., 1998), the primary language of the participants in this study. Using the recommended clinical cutoff of 13 (Cox et al., 1987), which has been used in prior studies in South Africa (Donald et al., 2019; Pellowski et al., 2023), 18.52% of the participants included in the main analysis at either time point had EPDS scores of 13 or greater (24.62% of participants in the group that endorsed alcohol use at any point in pregnancy met this cutoff and 14.43% of participants in the group that did not endorse alcohol use met this cutoff). The study validating the scale in Xhosa has a recommended cutoff of 12 or greater (Lawrie et al., 1998), and using this criterion, 20.37% of the participants included in the main analysis at either time point had EPDS scores of 12 or greater (29.23% of participants in the group that endorsed alcohol use at any point in pregnancy met this cutoff and 14.43% of participants in the group that did not endorse alcohol use met this cutoff).

All enrolled infants received a comprehensive medical exam at each visit, which included assessments of eye-related conditions. Several infants ($n = 3$) were identified as having eye-related anomalies during the medical exam, and they were excluded from any further analyses (note that none of these infants had reported PAE).

Substance Exposure Measurement

Alcohol use during pregnancy was measured by maternal self-report using the Alcohol Exposure Questionnaire (AEQ; Kwiatkowski et al., 2018) collected at enrollment. Due to the personal nature of these questions and learning from prior experience working with this population, these questions were asked as multiple-choice options of ordinal ranges. During this questionnaire, mothers were asked to estimate, on average, how many times they drank per week in each trimester (i.e., "Once per week or less," "2 to 3 times per week," "4 to 6 times per week," "Daily"). These responses were coded as the midpoint of the range unless the multiple choice option specified a specific number (i.e., 0.5 weekly drinking episodes, 2.5, 5, or 7, respectively). They were then asked an additional question about how many drinks were consumed per drinking


episode in that trimester (i.e., "Fewer than 2," "2 to 3," "4 or more," or "Not Sure"). Responses were coded as the midpoint of the range for the first two options (i.e., 1, 2.5, respectively). Those selecting "4 or more" were asked to write in a specific number which was used as the number of drinks per episode and those selecting "Not Sure" wrote in drink types such as "4 ciders" or "6 × 330 mL" which were later coded for alcohol units with a standardized system developed based on the Drakenstein study and the alcohol unit conversion scale from the South African Association for Alcohol Responsibility and Education (AWARE) and in consultation with an expert in PAE in South Africa (Stein et al., 2015; The Association for Alcohol Responsibility and Education, n.d.).

As our continuous measure for level of PAE, we extracted the number of standardized alcohol units consumed per week in each trimester from the AEQ (number of drinking episodes per week in each trimester × number of drinks consumed per episode). We also developed a continuous measure of exposure to binge drinking in each trimester (with a binge drinking episode defined as drinking 4 or more drinks in a particular episode) by extracting the number of drinking episodes per week for those who endorsed 4 or more drinks per episode in that trimester (CDC, 2022). See Figures 1 and 2 for frequency distributions of the number of drinks per week by trimester and of binge episodes per week by trimester, respectively, and Table 1 for descriptive statistics of alcohol endorsement.

One hundred three (26.2%) mothers endorsed alcohol use at any point in pregnancy out of the 393 mothers for whom that information was collected. Notably, for participants contributing to Visit 1 and/or Visit 2 data, there were no statistically significant differences between reported alcohol use patterns in mothers who endorsed alcohol use (without comorbid nonalcohol substance use) who were recruited prenatally compared to those recruited postnatally for drinks consumed per week in any trimester (Visit 1 sample: trimester 1: $p > .05$, $n = 52$; trimester 2 and trimester 3: $p > .05$, $n = 53$; Visit 2 sample: trimester 1, trimester 2, and trimester 3: $p > .05$, $n = 50$) or for binge episodes in trimester 1 (Visit 1 sample: $p > .05$, $n = 52$; Visit 2 sample: $p > .05$, $n = 50$), indicating consistent reporting across recruitment windows in those who endorsed alcohol use. There were also no statistically significant differences between reported alcohol use patterns as a function of pre- compared to postnatal recruitment in the overall sample of mothers whose infants participated in EEG visits (both those who endorsed and those who did not endorse prenatal alcohol use; without comorbid nonalcohol substance use) for any trimester (Visit 1 sample: trimester 1: $p > .05$, $n = 227$; trimester 2 and trimester 3: $p > .05$, $n = 228$; Visit 2 sample: trimester 1, trimester 2, and trimester 3: $p > .05$, $n = 226$) or for binge episodes in trimester 1 (Visit 1 sample: $p > .05$, $n = 227$; Visit 2 sample: $p > .05$, $n = 226$), suggesting consistent endorsement patterns in the overall sample regardless of recruitment timing.

In addition, alcohol risk scores were calculated using the WHO ASSIST (WHO ASSIST Working Group, 2002). This is a screening tool developed by a group of international researchers for use in identifying those most at risk of substance use problems in primary care settings (WHO ASSIST Working Group, 2002). To the best of our knowledge, there have not been any studies empirically evaluating its use in measuring alcohol risk in pregnant populations. In an effort to better predict which mothers would be at increased risk of alcohol use during pregnancy, we looked at whether the alcohol risk score calculated by the WHO ASSIST was related to reported drinking behavior and patterns across trimesters. We looked at correlations between the continuous alcohol risk score (measured


Figure 1
Number of Drinks per Week by Trimester (Nonwinsorized)




Note. These are frequency distributions of the sample ($n = 65$) with PAE, no comorbid nonalcohol substance exposure, and usable EEG data at either time point that is included in the main analyses. EEG = electroencephalography; PAE = prenatal alcohol exposure. See the online article for the color version of this figure.

Figure 2
Number of Binge Episodes per Week by Trimester (Nonwinsorized)


(a) Trimester 1



(b) Trimester 2



(c) Trimester 3



Note. These are frequency distributions of the sample ($n = 65$) with PAE, no comorbid nonalcohol substance exposure, and usable EEG data at either time point that is included in the main analyses. EEG = electroencephalography; PAE = prenatal alcohol exposure. See the online article for the color version of this figure.

Table 1

Descriptive Statistics of Number of Drinks per Week in Each Trimester and Number of Binge Episodes per Week in Each Trimester for Those Endorsing Alcohol Use at Any Point in Pregnancy (Nonwinsorized; $n = 65$)

Drinking variable	<i>n</i> -endorsing	<i>M</i>	<i>SD</i>	<i>Mdn</i>	Range	
					Min	Max
Number of drinks per week in						
Trimester 1	56	2.829	5.677	0.875	0.000	30.000
Trimester 2	29	1.390	4.506	0.000	0.000	30.000
Trimester 3	13	0.210	0.503	0.000	0.000	2.400
Number of binge episodes per week in						
Trimester 1	14	0.203	0.547	0.000	0.000	2.500
Trimester 2	6	0.077	0.333	0.000	0.000	2.500
Trimester 3	2	0.015	0.087	0.000	0.000	0.500

Note. These are descriptive statistics of the sample with PAE, no comorbid nonalcohol substance exposure, and usable EEG data at either time point that is included in the main analyses. Min = minimum; Max = maximum; EEG = electroencephalography; PAE = prenatal alcohol exposure.

from 0 to 39) and the number of drinks per week in each trimester and also the number of binge episodes per week in each trimester for the subsample of infants whose mothers completed both alcohol measures ($n = 392$; Figure 3).

The continuous alcohol risk score calculated using the WHO ASSIST significantly correlated with whether alcohol was dichotomously endorsed at any point during pregnancy, $r(390) = .44$, $p < .001$, as well as the continuous measure of number of drinks per week endorsed in each trimester, trimester 1: $r(390) = .31$, $p < .001$; trimester 2: $r(390) = .45$, $p < .001$; trimester 3: $r(390) = .22$, $p < .001$, and the continuous measure of number of binge episodes per week in each trimester, trimester 1: $r(390) = .21$, $p < .001$; trimester 2: $r(390) = .41$, $p < .001$; trimester 3: $r(390) = .21$, $p < .001$. These analyses indicate consistent reporting across measures of alcohol use behavior in the sample. Although only 16 mothers scored within the moderate- or high-risk WHO ASSIST categories, 100% of these mothers endorsed drinking during pregnancy, indicating the WHO ASSIST scores may be useful for identifying those who will drink during pregnancy to facilitate targeting counseling and other support services concerning the effects of PAE.

Non alcohol comorbid substance use endorsement in the 3 months prior to enrollment was collected using the WHO ASSIST (WHO ASSIST Working Group, 2002). Mothers reported on how often they endorsed tobacco, cannabis, cocaine, amphetamine, inhalant, sedative/sleeping pill, hallucinogen, opioid, or other drug use in the past three months prior to enrollment. Those who endorsed use of these nine substance categories in any amount or at any point in the past three months were coded as having comorbid substance use. Those who did not endorse use of any of these substances in the past three months were considered to have no comorbid substance use during pregnancy. An additional 21 of the 103 infants with PAE also had prenatal comorbid nonalcohol substance exposure and were excluded from further analyses for the purposes of this project.

Matching Comparison Participants for Primary Analyses

For each timepoint (i.e., Visit 1 or Visit 2), a matched subsample of infants without PAE or other substance exposure with usable EEG data were selected for comparison to the infants with PAE using the *MatchIt* package in R (Ho et al., 2011). These samples were matched

with *MatchIt* based on infant age at EEG data collection. At each timepoint, matched samples without PAE were not significantly different from those with exposure on the following potential confounding variables: infant age at EEG visit, maternal reports of depression as indexed by the EPDS (Cox et al., 1987), maternal age at infant birth, maternal education, and maternal income (Student's *t*-tests, all $p > .05$). Additionally, χ^2 tests indicated there was no significant difference in infant sex for those with alcohol exposure or without for each matched sample (both $p > .05$). See Figure 4 for a representation of how the final study sample across visits was determined. Demographic characteristics of the sample who contributed data at either time point are summarized in Table 2.


EEG Data Acquisition

EEG data were acquired at both Visit 1 and Visit 2 from infants while they were seated in their caregiver's lap in a dimly-lit, quiet room using a 128-channel high-density HydroCel Geodesic Sensor Net (EGI, Eugene, OR), amplified with a NetAmps 400 high-input amplifier, and recorded via an Electrical Geodesics, Inc. (EGI, Eugene, OR) system with a 1,000 Hz sampling rate. EEG data were online referenced to the vertex (channel Cz) through the EGI Netstation software. Impedances were kept below 100 k Ω in accordance with the impedance capabilities of the high-impedance amplifiers. Geodesic Sensor Nets with modified tall pedestals designed for improving inclusion of infants with thick/curly/tall hair were used as needed across participants. Shea moisture leave-in castor oil conditioner was applied to hair across the scalp prior to net placement to improve both impedances and participant comfort. This leave-in conditioner contains insulating ingredients so there is no risk of electrical bridging and has not been found to disrupt the EEG signal during testing (unpublished data). Conditioning hair in this way allows for nets to lay closer to the scalp for Afro-textured hair types and makes for far more comfortable net removal at the end of testing.

The VEP task was presented using Eprime 3.0 software (Psychology Software Tools, Pittsburgh, PA) on a Lenovo desktop computer with an external monitor 19.5 in. on the diagonal facing the infant (with monitor approximately 65 cm away from the infant). A standard phase-reversal VEP was induced with a black and white checkerboard (1 cm \times 1 cm squares within the board) stimulus that alternated presentation (black squares became white, white squares became black) every 500 ms for a total of 100 trials. Participant looking was monitored by

Figure 3

WHO ASSIST Alcohol Risk Scores and Number of Drinks per Week by Trimester (a–c) and Number of Binge Episodes per Week by Trimester (d–f)



Note. Sample size ($n = 392$). Axes vary by trimester to best capture the range of alcohol endorsed in that trimester. WHO ASSIST = World Health Organization Alcohol, Smoking and Substance Involvement Screening Test. See the online article for the color version of this figure.

video and by an assistant throughout data collection. If the participant looked away during the VEP task, the task was rerun.


EEG Data Preprocessing

VEP data were exported from native Netstation .mff format to .raw format and then preprocessed using the HAPPE + ER pipeline


within the HAPPE v3.3 software, an automated open-source EEG processing software validated for infant data (Monachino et al., 2022). A subset of the 128 channels were selected for preprocessing that excluded the rim electrodes as these are typically artifact-laden (channels excluded from preprocessing included in Table 3). The HAPPE + ER pipeline was run with user-selected specifications that are outlined in Table 3.

Figure 4
Participants Contributing EEG Data at Each Visit

(a) Visit 1 Participants (n=108)



(b) Visit 2 Participants (n=102)



Note. EEG = electroencephalography; VEP = visual-evoked potential; ERP = event-related potential. See the online article for the color version of this figure.

Table 2
Demographic Characteristics

Demographic characteristic	Total (N = 162)
Maternal place of birth	
South Africa	158 (97.5%)
In the African Continent (not South Africa)	4 (2.5%)
Primary spoken language	
Xhosa language	155 (95.7%)
English language	3 (1.9%)
Sotho language	2 (1.2%)
Afrikaans language	1 (0.6%)
Zulu language	1 (0.6%)
Maternal age at infant birth (years)	
M (SD)	28.9 (5.88)
Mdn (Min, Max)	29.0 (18.0, 43.0)
Missing	1 (0.6%)
Maternal educational attainment ^a	
Completed Grade 6 (Standard 4) to Grade 7 (Standard 5)	3 (1.9%)
Completed Grade 8 (Standard 6) to Grade 11 (Standard 9)	70 (43.2%)
Completed Grade 12 (Standard 10; i.e., completed high school)	63 (38.9%)
Some university/college/postmatric education	20 (12.3%)
Completed university/college/postmatric education	6 (3.7%)
Maternal monthly income (South African Rand—ZAR) ^b	
Less than R1,000 per month	90 (55.6%)
R1,000–R5,000 per month	53 (32.7%)
R5,000–R10,000 per month	13 (8.0%)
More than R10,000 per month	1 (0.6%)
Unknown	5 (3.1%)
EPDS depression score ^c	
M (SD)	6.85 (5.75)
Mdn (Min, Max)	6.00 (0, 22.0)
WHO ASSIST alcohol risk score ^d	
M (SD)	1.23 (4.53)
Mdn (Min, Max)	0 (0, 30.0)
Infant biological sex	
Female	78 (48.1%)
Male	82 (50.6%)
Missing	2 (1.2%)

Note. EPDS = Edinburgh Postnatal Depression Scale; WHO ASSIST = World Health Organization Alcohol, Smoking, and Substance Involvement Screening Test; Min = minimum; Max = maximum.

^aThe South African Educational System was formerly divided into years called standards, similar to the way the United States Educational System is divided into grades. The equivalent in terms of standards is provided in parentheses next to each mentioned grade. “University/college/postmatric education” refers to tertiary or postsecondary education as defined by the World Bank. ^bAt the time of writing (May 17, 2023), 1 USD = 19.24 South African Rand (ZAR). ^cDepression was measured using the EPDS at enrollment (Cox et al., 1987). ^dAlcohol risk score was calculated using the WHO ASSIST at enrollment (WHO ASSIST Working Group, 2002).

Preprocessed VEP data were considered usable and moved forward to VEP extraction if HAPPE preprocessing ran successfully, at least 15 trials were retained following bad trial rejection, and at least one good channel was kept within the visual region of interest (ROI). Note that channels marked bad during preprocessing had their data interpolated as part of standard preprocessing pipelines for ERPs (Monachino et al., 2022). Interpolated channels were included in analyses here as is typically done in developmental samples and given the low overall rates of interpolation present (e.g., all groups at all visits had an average of between 4 to 5 of 5 possible good channels in the ROI retained). At this point, 6 infants with PAE were excluded from further analysis from the Visit 1 time point, and 1 infant with PAE was excluded from the Visit 2 time point. Student’s *t*-tests confirmed that there were no significant differences in any data quality measure between the alcohol-exposed and nonexposed groups at either time point (all $p > .05$) with the exception of number and

percent of retained channels in the ROI for the alcohol-exposed versus nonexposed groups at Visit 2, (both $t(100) = 2.37$, $p = .020$). Regression models for Visit 2 were run including the number of useable channels within the ROI as a covariate. However, since this variable was not a significant predictor of any VEP outcome measure (all predictors $p > .05$) nor did it change any of the other model and/or predictors’ significance or any pattern of results in these analyses, it was removed from the final regression models reported in the results section to preserve model parsimony and statistical power. All EEG data quality metrics for included files at each age for each group (PAE or no exposure) are summarized in Table 4.

VEPs

VEP waveforms were extracted and quantified using the HAPPE v3.3 GenerateERPs script (Monachino et al., 2022). Electrodes in

Table 3
HAPPE v3.3 Preprocessing Script Parameters

Preprocessing parameter type	Parameter selection
Density	High (>30 channels)
Resting state or task	Task
ERP analysis	Yes
Acquisition layout	128-channel EGI HydroCel Geodesic Sensor Net
Channels	All except E1, E8, E14, E17, E21, E25, E32, E38, E43, E44, E48, E49, E56, E63, E68, E73, E81, E88, E94, E99, E107, E113, E114, E119, E120, E121, E125, E126, E127, E128
Line noise	
Line noise frequency	50 Hz
Line noise reduction method	CleanLine—default
Resample	Off
Filter	
Filter—low-pass cutoff	30 Hz
Filter—high-pass cutoff	0.3 Hz
Filter type	EEGLAB's FIR
Bad channel detection	On
Bad channel detection method	Default
Wavelet thresholding	Default
Wavelet threshold rule	Hard
Muscle	Off
Segmentation	On
Starting parameter for stimulus	−0.1 s
Ending parameter for stimulus	0.5 s
Task offset	11 ms
Baseline correction	On
Baseline correction start	−100 ms
Baseline correction end	0 ms
Interpolation	Off
Segment rejection	On
Segment rejection method	Amplitude criteria only
Minimum segment rejection threshold	−200
Maximum segment rejection threshold	200
Segment rejection based on all channels or ROI	ROI
ROI channels	E70, E71, E75, E76, E83
Rereference method	Average

Note. ERP = event-related potential; EGI = Electrical Geodesics, Inc.; FIR = finite impulse response filter; ROI = region of interest.

the occipital region were selected as a ROI (i.e., E70, E71, E75, E76, E83). The VEP waveform has three main components to be quantified: a negative N1 peak, a positive P1 peak, and a negative N2 peak. The feature of interest for these components was peak latency. The window for calculating peak latencies for the N1 component was 40–100 ms, 75–175 ms for the P1 component, and 100–325 ms for the N2 component. HAPPE parameters used in extracting the ERPs are summarized in Table 5.

All VEPs were visually inspected to ensure that the automatically extracted values were correct and were adjusted if observable peaks occurred outside the automated window bounds. Participants were considered to have failed this visual inspection and were subsequently removed from the data set if their VEP did not produce three discernible peaks.

Of the alcohol-exposed group, no infants from Visit 1 failed visual inspection, and seven infants from Visit 2 failed visual inspection. This left a final sample of 54 infants with PAE and no comorbid substance exposure with usable EEG data at Visit 1, and 51 infants with these exposure characteristics with usable EEG data at Visit 2 (40 of these infants also provided usable EEG data at Visit 1). The determination process for usable EEG is detailed in Figure 4. Data quality for included files at each age for each group (PAE or no exposure)

are summarized in Table 4. See Figure 5 for visualizations of the average VEP ERP waveforms at both visit time points.

Sample Included in VEP Analyses

After exclusions during processing, the final sample size for Visit 1 was 108 infants (54 with PAE, 54 matched comparisons with no prenatal substance exposure). There were no statistically significant differences between infants with PAE who contributed usable VEP and those who did not (either because VEP was not collected or because it was deemed unusable) with respect to maternal EPDS scores, education level, income, maternal age at birth (Student's *t*-tests, all $p > .05$), or infant sex (Pearson's χ^2 test, $p > .05$). Through the matching process described above, the comparison infants without PAE included in analyses from the larger pool of data were not statistically different from the infants with PAE on any of these variables, either.

The final sample size for Visit 2 was 102 infants (51 with PAE, 51 matched comparisons with no prenatal substance exposure). Again, there were no statistically significant differences between infants with PAE who contributed usable VEP and those who did not (either because VEP was not collected or because it was deemed unusable) with respect to maternal EPDS scores, income, age at birth, or infant sex (all $p > .05$).

Table 4
EEG Quality Control Measures

Quality control measure	Visit 1 data (<i>n</i> = 108)			Visit 2 data (<i>N</i> = 102)		
	Alcohol exposure (<i>n</i> = 54)	No alcohol exposure (<i>n</i> = 54)	<i>t</i> test between groups (α = .05)	Alcohol exposure (<i>n</i> = 51)	No alcohol exposure (<i>n</i> = 51)	<i>t</i> test between groups (α = .05)
VEP trial retention						
Number of collected VEP trials						
<i>M</i> (SD)	99.9 (0.408)	100 (0)	n.s.	100 (0)	100 (0)	n.s.
<i>Mdn</i> (Min, Max)	100 (97.0, 100)	100 (100, 100)		100 (100, 100)	100 (100, 100)	
Number of retained VEP trials						
<i>M</i> (SD)	92.9 (16.4)	92.5 (15.7)	n.s.	95.3 (13.7)	95.5 (12.8)	n.s.
<i>Mdn</i> (Min, Max)	100 (17.0, 100)	100 (32.0, 100)		100 (21.0, 100)	100 (29.0, 100)	
Percent retained VEP trials						
<i>M</i> (SD)	93.0 (16.5)	92.5 (15.7)	n.s.	95.3 (13.7)	95.5 (12.8)	n.s.
<i>Mdn</i> (Min, Max)	100 (17.0, 100)	100 (32.0, 100)		100 (21.0, 100)	100 (29.0, 100)	
Whole head channel retention						
Number of retained channels						
<i>M</i> (SD)	78.5 (8.83)	79.1 (9.37)	n.s.	78.7 (10.7)	81.5 (8.34)	n.s.
<i>Mdn</i> (Min, Max)	79.5 (56.0, 95.0)	78.5 (55.0, 97.0)		81.0 (46.0, 96.0)	83.0 (59.0, 96.0)	
Percent retained channels						
<i>M</i> (SD)	79.3 (8.92)	79.9 (9.47)	n.s.	78.7 (10.7)	81.9 (8.18)	n.s.
<i>Mdn</i> (Min, Max)	80.3 (56.6, 96.0)	79.3 (55.6, 98.0)		80.8 (46.5, 94.9)	82.8 (59.6, 93.9)	
Region of interest channel retention						
Number of retained channels						
<i>M</i> (SD)	4.24 (0.930)	4.31 (1.01)	n.s.	4.04 (1.06)	4.47 (0.758)	<i>t</i> (100) = 2.37
<i>Mdn</i> (Min, Max)	4.50 (2.00, 5.00)	5.00 (1.00, 5.00)		4.00 (1.00, 5.00)	5.00 (2.00, 5.00)	<i>p</i> = .020
Percent retained channels						
<i>M</i> (SD)	0.848 (0.186)	0.863 (0.201)	n.s.	0.808 (0.212)	0.894 (0.152)	<i>t</i> (100) = 2.37
<i>Mdn</i> (Min, Max)	0.900 (0.400, 1.00)	1.00 (0.200, 1.00)		0.800 (0.200, 1.00)	1.00 (0.400, 1.00)	<i>p</i> = .020
Correlation of data pre- versus postwavelet thresholding (Pearson's <i>r</i>)						
At 5 Hz						
<i>M</i> (SD)	0.529 (0.192)	0.500 (0.184)	n.s.	0.455 (0.226)	0.384 (0.251)	n.s.
<i>Mdn</i> (Min, Max)	0.569 (0.0838, 0.901)	0.514 (0.0389, 0.835)		0.484 (0.0497, 0.795)	0.401 (0.0317, 0.818)	
At 8 Hz						
<i>M</i> (SD)	0.450 (0.176)	0.411 (0.165)	n.s.	0.425 (0.235)	0.354 (0.222)	n.s.
<i>Mdn</i> (Min, Max)	0.459 (0.0884, 0.844)	0.421 (0.0689, 0.763)		0.440 (0.0240, 0.805)	0.349 (0.0262, 0.764)	
At 12 Hz						
<i>M</i> (SD)	0.428 (0.166)	0.405 (0.159)	n.s.	0.383 (0.237)	0.300 (0.193)	n.s.
<i>Mdn</i> (Min, Max)	0.439 (0.0702, 0.798)	0.442 (0.0377, 0.644)		0.400 (0.0179, 0.762)	0.323 (0.0159, 0.623)	
At 20 Hz						
<i>M</i> (SD)	0.492 (0.177)	0.481 (0.164)	n.s.	0.431 (0.249)	0.358 (0.220)	n.s.
<i>Mdn</i> (Min, Max)	0.533 (0.100, 0.832)	0.506 (0.0839, 0.757)		0.475 (0.0229, 0.830)	0.399 (0.0145, 0.737)	

Note. EEG data were preprocessed and VEPs were extracted using HAPPE + ER v3.3 software, an automated open-source EEG processing software validated for infant data (Monachino et al., 2022). EEG = electroencephalography; VEP = visual-evoked potential; Min = minimum; Max = maximum; n.s. = nonsignificant.

Table 5
HAPPE v3.3 GenerateERPs Script Parameters

GenerateERPs parameter type	Parameter selection
Average or individual trials	Average
Channels of interest	E70, E71, E75, E76, E83
Bad channels included/excluded	Included
Calculating ERP values	On
Windows	Min 40–100 ms Max 75–175 ms Min 100–325 ms

Note. ERP = event-related potential; Min = minimum; Max = maximum.

Infants with PAE included in VEP analyses had significantly higher maternal education levels than those that did not contribute, $t(80) = -2.45$, $p = .017$, though both groups had mean education levels corresponding to the same rank of schooling completion (i.e., Standard 6 [Grade 8] to Standard 9 [Grade 11]). Through matching, the comparison of infants without PAE was not statistically different from the infants with PAE on any of these variables.

Participants Included in Exploratory MRI Analyses

To better explore the potential of myelination as a mechanism explaining VEP latencies with limited MRI data, participants were not split into Visit 1 or Visit 2 groups for these analyses. Instead, our sample was composed of infants who had usable data for both EEG and MRI at either time point. Thirty-four infants had usable EEG and MRI data from the Visit 1 time point, and 14 infants had usable EEG and MRI data from the Visit 2 time point. Only one infant had usable EEG and MRI data at both of their visits, so we selected one of their visits (Visit 2) for inclusion in this analysis (note that results remained the same regardless of

which visit was included for this individual, if both visits were included, or if neither was included; final $n = 47$ measurements modeled). In the sample for this exploratory analysis, 19.15% of infants ($n = 9$) had PAE, while the remainder (80.85%, $n = 38$) had no PAE. Infants with usable neuroimaging data were included in the sample regardless of prenatal substance exposure (i.e., alcohol or no alcohol exposure) as the sample was not powered to explicitly explore the impact of PAE on this relation.


MRI Data Acquisition

MRI data including T1- and T2-weighted scans were collected during natural sleep on a 3T Siemens MAGNETOM Skyra scanner (Siemens Medical Solutions USA, Inc., Malvern, PA) with a 16-channel neonatal head coil. Infants were scheduled for MRI scans during their normal nap time whenever possible. Before scanning, infants were fed and then swaddled according to the feed and wrap technique and then soothed to sleep by their caregiver (Antonov et al., 2017). Ear cushions and headphones were worn by infants to adequately protect their hearing. If the infant woke up or was distressed at any point, scanning would immediately stop and start again once the infant was settled and if the caregiver wished to continue. Each participant had structural imaging, acquired with the following parameters: T₁ MPRAGE (echo time [TE] = 1.69 ms, repetition time [TR] = 2,400 ms, 1 mm isotropic voxel, 256 × 256 mm field of view, inversion time [TI] = 1,450 ms, flip angle = 7°, 4 min 10 s scanning time); T₂ (TE = 561 ms, TR = 3,200 ms, TI = 1,800 ms, 1 mm isotropic voxel, 250 × 250 mm field, flip angle = 120° of view, 6 min 35 s scanning time).

T1w/T2w Ratios

T1w/T2w ratio mapping was performed to generate images with contrast related to myelin content (Ganzetti et al., 2014; Glasser &

Figure 5
Average Visual-Evoked Potential Waveforms at Each Visit



Note. VEP = visual-evoked potential. See the online article for the color version of this figure.

Van Essen, 2011). This workflow was conducted using the MRtool tool box (Ganzetti et al., 2014, 2015, 2018) in SPM8 (Wellcome Trust Centre for Neuroimaging, London, United Kingdom) with Matlab 2023a (Mathworks).

Initially, T2w images were coregistered to T1w, followed by separate intensity normalization and segmented into tissue classes with “New Segment” in SPM (including gray matter white matter and cerebrospinal fluid). Intensity normalization corrects for bias introduced due to slow wave variation and individual cranial structure. Further intensity scaling for both T1w and T2w is calculated by using masked regions outside of the brain to ensure intensity scaling is comparable between subjects (Ganzetti et al., 2014). Following this calibration, images were transformed into standard Montreal Neurological Institute (MNI) space for further statistical analysis. Each participant’s image was visually inspected for quality assurance. ROI masks were generated from the Glasser et al. (2016) atlas and registered to standard space. T1w/T2w ratio estimates were calculated for ROIs, specifically a priori regions for statistical analysis (i.e., V1, MT/V5).

Statistical Analyses

All analyses were performed using R Statistical Software (v4.2.1; R Core Team, 2022) in the RStudio interface (v2022.7.0.548; RStudio Team, 2015), and multiple regression analyses were performed using the *stats* package in R (R Core Team, 2022). For each analysis, overly influential data points in each regression model were tested for using the *MASS* package in R (Venables & Ripley, 2003) by evaluating studentized residuals with a priori cutoff bounds of -3 and 3 residual deviations. Any detected outliers were removed from the models, which were then rerun. R scripts for regression analyses are available on the Open Science Framework (Margolis, 2024).

VEP Analyses

Given the presence of several extreme scores in the distributions of PAE variables and the potential for these scores to overly influence the subsequent regression models, we elected to winsorize drinking scores using the *DescTools* package in R (Signorelli, 2023). Within each visit timepoint, for those endorsing alcohol at any point in their pregnancy, scores above the upper 95% quantile of number of drinks endorsed per week in the first, second, and third trimester as well as number of binge episodes per week in the first trimester were winsorized separately (see Figures S1–S4 in the online supplemental materials for frequency distributions of the winsorized number of drinks per week by trimester and of winsorized binge episodes in the first trimester and Table S1 in the online supplemental materials for descriptive statistics of winsorized alcohol endorsement values). Specifically, these extreme scores were given the same value as the score associated with the 95% quantile for that trimester and retained in analyses. This strategy facilitated preserving model parsimony (modeling multiple trimesters simultaneously) and sample size and statistical power (without needing to remove all participants with extreme values in any of the trimesters modeled).

Maternal age at infant birth, maternal education, maternal income, and EPDS depression scores were considered as potential confounding covariates of drinking scores. In both the Visit 1 ($n = 108$) and Visit 2 samples ($n = 102$), maternal age at infant birth, maternal

education, maternal income, and EPDS depression scores were not significantly related to winsorized drinking variables for any trimester used as model predictors (i.e., number of drinks per week in each trimester and number of binge episodes per week in trimester 1; all $p > .05$).

To correct for the potential influence of earlier component latency differences on later component latencies, corrected latencies were calculated and used in all VEP latency analyses of alcohol exposure. Specifically, the P1 latency was corrected for the N1 latency (corrected P1 latency = $P1 - N1$ latency), and the N2 latency was corrected for the P1 latency (corrected N2 latency = $N2 - P1$ latency).

Three ordinary-least squares multiple regression models were run for each visit timepoint, one for each VEP component latency as the outcome (i.e. N1, P1, N2 latencies). Models were run including number of drinks per week for each trimester (three predictors total, one per trimester). Infant age at data collection and number of retained usable VEP trials were included as a priori covariates in all models. To test whether nonlinear age-related effects should also be included in regression models during each visit age range, model comparisons were performed on models with (a) linear age effects, (b) linear and natural log(age) effects, and (c) linear and quadratic age effects using the *AICcmodavg* and *stats* package in R (Mazerolle, 2023; R Core Team, 2022). Only the Visit 1 P1 latency model indicated a statistically significantly better fit with the inclusion of a nonlinear age term (Visit 1 P1: $F(1, 3) = 27.83$, $p = .0000008$; all other tests: $p > .05$) so Visit 1 P1 results are reported including both age and natural log(age) as predictors in the model (though note the pattern of results with respect to drinking predictors remains the same with and without the nonlinear age term).

Multivariate outliers in each regression model were evaluated using the *MASS* package in R (Venables & Ripley, 2003). We considered data points to be overly influential by evaluating studentized residuals with a priori cutoff bounds of -3 and 3 residual deviations. Multivariate outliers were removed, and analyses were rerun if any such outliers were detected for each model. No more than three participants were removed from each model of number of drinks per week using this criterion. Note that one participant with Visit 1 data was missing a continuous measure of alcohol exposure in the first trimester and was therefore removed from relevant models for missingness.

To control for the number of model comparisons per question (three models, one for each VEP component), Bonferroni correction was applied to each set of models ($\alpha = .05$, corrected p threshold per model = $.016$). Significant predictors were examined only in models that survived correction for multiple comparisons.

To examine the effects of binge-drinking behavior, three additional ordinary-least squares regression models were run for each visit timepoint, one for each VEP component latency as the outcome (i.e. N1, P1, N2 latencies). Models were run including number of binge episodes per week for the first trimester as a predictor. Infant age at data collection and number of retained usable VEP trials were included as a priori covariates in all models. Nonlinear age terms were tested for potential inclusion as above, and again, only the Visit 1 P1 regression model indicated significantly better fit with the inclusion of a nonlinear age term (P1: $F(1, 3) = 27.86$, $p = .0000007$; all other tests: $p > .05$) so Visit 1 P1 results are reported including both age and natural log(age) as predictors (though note the pattern of results with respect to drinking predictors remains the same with and without the nonlinear age term). No more than three participants were removed from each model of number of binge episodes per week using the criterion for outliers described

above. To control for the number of model comparisons per question at each visit (three models, one for each VEP component), Bonferroni correction was applied to each set of models ($\alpha = .05$, corrected p threshold per model = .016). Significant predictors were examined only in models that survived correction for multiple comparisons.

Over the two sets of analyses of interest, (a) effects of drinks per week over three trimesters and (b) effects of binge episodes per week in the first trimester examined at each of the two visit timepoints, a total of 12 regression models were run.

Exploratory Analyses of T1w/T2w Ratios in Relation to VEP Latencies

To explore a possible postnatal influence on VEP latencies, we looked at the effect of myelination in the visual cortex on VEP latencies in a cross-sectional subsample of participants. To create this subsample, we collated all participants who had usable VEP data and T1w/T2w ratios at the same time point (i.e., both EEG and MRI data at Visit 1 or Visit 2). We selected myelination in the primary visual cortex (V1) and in the middle temporal visual area (i.e., MT or V5) as our brain ROIs based on prior work with adults (Barnikol et al., 2006). For both ROI, we averaged estimates of myelination lateralized across the left and right hemispheres to create an average value for each ROI. Here we examined VEP latencies without correction as multiple components are derived from the visual regions tested (e.g., correcting the latencies for later components would remove the shared variance of interest for this analysis). We ran a multiple regression looking at the impact of myelination in both of these ROI (as indexed by the T1w/T2w ratio) on each component VEP latency (running six models total) with infant age at EEG collection, number of retained VEP trials, and infant age at MRI collection as covariates. No more than two participants were removed from each model of myelination using the criterion for outliers described above. No multiple comparison correction was applied given the exploratory nature of the analyses, though notably, the significant effect observed would survive such correction. Due to the smaller sample sizes of this exploratory analysis (only 19.15% of infants had PAE), we did not have power to explore the impact of alcohol exposure on this relation directly.

Results

Effects of Number of Drinks Per Week on VEP Latencies

At Visit 1 (Ages 8.286–24.571 Weeks)

In the Visit 1 subsample (age in weeks: $M = 15.798$, $SD = 3.719$, range = 8.286–24.571, total n before removing model outliers = 108), the ordinary-least squares regression model examining the effect of number of drinks per week in each trimester, controlling for infant age at Visit 1 EEG collection, the natural log(age at Visit 1), and number of retained VEP trials, significantly explained variance in the corrected P1 latency ($n = 104$, model $R^2 = .56$, $F(6, 97) = 20.89$, $p < .001$) and survived correction for multiple comparisons across models (corrected threshold $p = .016$). Within the model, number of drinks per week in the first trimester (controlling for number of drinks in the second and third trimesters) significantly predicted corrected P1 latency ($b = 1.16$, $p = .039$, 95% confidence interval [CI] [0.06, 2.26]), such that those with increased alcohol exposure had longer corrected P1 latencies (i.e. P1–N1 latency; see Figure 6 for residual plot). In this model, infant age at Visit 1 EEG collection and the natural log(age)

also significantly predicted corrected P1 latency over this age range (age: $b = 7.36$, $p < .001$, 95% CI [3.61, 11.11]; natural log(age): $b = -153.50$, $p < .001$, 95% CI [-212.39, -94.61]) such that increasing age was associated with shorter P1 latencies.

Neither the N1 nor the N2 VEP components showed statistically significant effects of drinks per week in any trimester over this age range. The model examining effects on the N1 latency was not statistically significant ($n = 105$, model $R^2 = .08$, $F(5, 99) = 1.81$, $p = .117$). The model examining effects on the corrected N2 latency was statistically significant ($n = 106$, model $R^2 = .25$, $F(5, 100) = 6.60$, $p < .001$) and survived correction for multiple comparisons (corrected threshold $p = .016$). However, number of drinks per week in any trimester did not significantly predict corrected N2 latency (all trimesters $p > .05$). See Table 6 for a regression table of number of drinks per week in each trimester predicting each VEP component at Visit 1.

At Visit 2 (Ages 26.000–51.857 Weeks)

In the Visit 2 subsample (age in weeks: $M = 36.489$, $SD = 6.337$, range = 26.000–51.857, total n before removing model outliers = 102), there were no statistically significant effects of drinks per week in any trimester on any VEP component latency (N1, P1, N2). Specifically, the model examining the effect of number of drinks per week in each trimester significantly explained variance in the N1 latency, controlling for infant age at Visit 2 EEG collection and number of retained VEP trials, but did not survive correction for multiple comparisons across models ($n = 102$, model $R^2 = .11$, $F(5, 96) = 2.46$, $p = .038$; corrected threshold $p = .016$). Neither P1 nor N2 latency models were statistically significant (P1: $n = 100$, model $R^2 = .03$, $F(5, 94) = 0.64$, $p = .671$; N2: $n = 101$, model $R^2 = .02$, $F(5, 95) = 0.34$, $p = .889$). See Table 7 for a regression table of number of drinks per week in each trimester predicting each VEP component at Visit 2.

Effects of Binge Drinking Behaviors on VEP Latencies


When considering the effect of drinking pattern (i.e., binge drinking behaviors), sufficient variability to model effects existed only for trimester 1 (trimester 1 mothers reporting binge drinking $n = 14$) but not subsequent trimesters (trimester 2 mothers reporting binge drinking $n = 6$, trimester 3 mothers reporting binge drinking $n = 2$).

At Visit 1 (Ages 8.286–24.571 Weeks)

In the Visit 1 subsample (total n before removing model outliers = 107), the ordinary-least squares regression model examining the effect of number of binge episodes per week in the first trimester, controlling for infant age at Visit 1 EEG collection, natural log(age at Visit 1) and number of retained VEP trials, significantly explained variance in the corrected P1 latency ($n = 104$, model $R^2 = .56$, $F(4, 99) = 30.99$, $p < .001$) and survived correction for multiple comparisons across models (corrected threshold $p = .016$). Within the model, number of binge episodes in trimester 1 significantly predicted corrected P1 latency ($b = 9.14$, $p = .028$, 95% CI [0.98, 17.31]), such that those with increased binge exposure had longer P1 latencies (see Figure 7 for residual plot). In this model, infant age at Visit 1 EEG collection and the natural log(age) also significantly predicted corrected P1 latency over this age range (age: $b = 7.47$, $p < .001$, 95% CI [3.72, 11.21]; natural log[age]:

Figure 6

Residual Plot for Number of Drinks per Week in Trimester 1 Predicting Corrected P1 Latency at Visit 1 (n = 104)



Note. See the online article for the color version of this figure.

$b = -154.65$, $p < .001$, 95% CI $[-213.41, -95.89]$ such that increasing age was associated with shorter P1 latencies.

Neither the N1 nor the N2 VEP components showed statistically significant effects of binge episodes per week over this age range. The model examining effects on the N1 latency was statistically significant but did not survive correction for multiple comparisons (corrected threshold $p = .016$; $n = 105$, model $R^2 = .07$, $F(3, 101) = 2.71$,

$p = .049$). The model examining effects on the corrected N2 latency was statistically significant ($n = 106$, model $R^2 = .24$, $F(3, 102) = 10.92$, $p < .001$) and survived correction for multiple comparisons. However, number of binge episodes per week in the first trimester did not significantly predict corrected N2 latency. See Table 8 for a regression table of number of binge episodes per week in the first trimester predicting each VEP component at Visit 1.

Table 6

Regression Models of Number of Drinks per Week in Each Trimester Predicting VEP Component Latencies at Visit 1 (n Before Removing Outliers = 108)

Component	Variable	<i>b</i>	<i>SE</i>	<i>p</i>	95% CI [LL, UL]
N1 latency	Model: $n = 105$, model $R^2 = .08$, $F(5, 99) = 1.81$, $p = .117$				
	Number of drinks per week in				
	Trimester 1	−0.23	0.49	.646	[−1.20, 0.74]
	Trimester 2	−2.21	1.39	.116	[−4.96, 0.55]
	Trimester 3	−0.21	2.89	.941	[−5.96, 5.53]
	Infant age at EEG collection	−0.13	0.23	.566	[−0.60, 0.33]
P1 latency	Model: $n = 104$, model $R^2 = .56$, $F(6, 97) = 20.89$, $p < .001$				
	Number of drinks per week in				
	Trimester 1	1.16	0.55	.039	[0.06, 2.26]
	Trimester 2	0.18	1.53	.906	[−2.86, 3.22]
	Trimester 3	−5.04	3.23	.121	[−11.45, 1.36]
	Infant age at EEG collection	7.36	1.89	<.001	[3.61, 11.11]
N2 latency	Natural log of infant EEG age	−153.50	29.67	<.001	[−212.39, −94.61]
	Number of retained EEG trials	0.09	0.06	.142	[−0.03, 0.20]
	Model: $n = 106$, model $R^2 = .25$, $F(5, 100) = 6.60$, $p < .001$				
	Number of drinks per week in				
	Trimester 1	2.92	2.00	.148	[−1.05, 6.90]
	Trimester 2	−2.00	5.70	.726	[−13.31, 9.31]
	Trimester 3	0.83	11.86	.944	[−22.70, 24.36]
	Infant age at EEG collection	−5.17	0.93	<.001	[−7.02, −3.32]
	Number of retained EEG trials	0.14	0.22	.514	[−0.29, 0.57]

Note. VEP = visual-evoked potential; CI = confidence interval; LL = lower limit; UL = upper limit; EEG = electroencephalography.

Table 7

Regression Models of Number of Drinks per Week in Each Trimester Predicting VEP Component Latencies at Visit 2 (n Before Removing Outliers = 102)

Component	Variable	<i>b</i>	<i>SE</i>	<i>p</i>	95% CI [LL, UL]
N1 latency	Model: $n = 102$, model $R^2 = .11$, $F(5, 96) = 2.46$, $p = .038$				
	Number of drinks per week in				
	Trimester 1	0.21	0.20	.281	[-0.18, 0.60]
	Trimester 2	0.23	0.54	.669	[-0.84, 1.31]
	Trimester 3	0.51	2.10	.810	[-3.66, 4.67]
	Infant age at EEG collection	-0.28	0.09	.002	[-0.46, -0.10]
P1 latency	Number of retained EEG trials	0.01	0.04	.804	[-0.08, 0.10]
	Model: $n = 100$, model $R^2 = .03$, $F(5, 94) = 0.64$, $p = .671$				
	Number of drinks per week in				
	Trimester 1	0.06	0.15	.689	[-0.24, 0.37]
	Trimester 2	-0.54	0.43	.207	[-1.39, 0.31]
	Trimester 3	-0.14	1.66	.931	[-3.43, 3.14]
N2 latency	Infant age at EEG collection	-0.06	0.07	.378	[-0.20, 0.08]
	Number of retained EEG trials	-0.02	0.03	.566	[-0.09, 0.05]
	Model: $n = 101$, model $R^2 = .02$, $F(5, 95) = 0.34$, $p = .889$				
	Number of drinks per week in				
	Trimester 1	0.09	0.58	.874	[-1.06, 1.24]
	Trimester 2	-1.58	1.60	.326	[-4.75, 1.59]
	Trimester 3	-3.57	6.92	.607	[-17.31, 10.17]
	Infant age at EEG collection	-0.17	0.27	.515	[-0.70, 0.35]
	Number of retained EEG trials	0.06	0.13	.650	[-0.20, 0.31]

Note. VEP = visual-evoked potential; CI = confidence interval; LL = lower limit; UL = upper limit; EEG = electroencephalography.


At Visit 2 (Ages 26.000–51.857 Weeks)

In the Visit 2 subsample (total n before removing model outliers = 102), there were no statistically significant effects of binge episodes per week in the first trimester on any VEP component latency (N1, P1, N2). Specifically, the model examining the effect of number of binge episodes per week in the first trimester on the N1 latency, controlling for infant age at Visit 2 EEG collection and number of retained VEP trials, was statistically significant ($n = 101$, model $R^2 = .11$,

$F(3, 97) = 3.94$, $p = .011$) and survived correction for multiple comparisons, but the number of binge episodes in the first trimester did not significantly predict N1 latency ($p > .05$). The model examining the effect of number of binge episodes per week in the first trimester did not significantly explain variance in the corrected P1 latency ($n = 100$, model $R^2 = .02$, $F(3, 96) = 0.56$, $p = .645$). The model examining the effect of number of binge episodes per week in the first trimester did not significantly explain variance in the corrected N2 latency either ($n = 101$, model $R^2 = .01$, $F(3, 97) = 0.47$, $p = .702$). See

Figure 7

Residual Plot for Number of Binge Episodes per Week in Trimester 1 Predicting Corrected P1 Latency at Visit 1 (n = 104)



Note. See the online article for the color version of this figure.

Table 8

Regression Models of Number of Binge Episodes per Week in Each Trimester Predicting VEP Component Latencies at Visit 1 (n Before Removing Outliers = 108)

Component	Variable	<i>b</i>	<i>SE</i>	<i>p</i>	95% CI [LL, UL]
N1 latency	Model: $n = 105$, model $R^2 = .07$, $F(3, 101) = 2.71$, $p = .049$				
	Number of binge episodes per week in Trimester 1	-5.93	3.34	.079	[-12.56, 0.70]
	Infant age at EEG collection	-0.09	0.23	.710	[-0.54, 0.37]
	Number of retained EEG trials	-0.11	0.05	.040	[-0.21, 0.00]
P1 latency	Model: $n = 104$, model $R^2 = .56$, $F(4, 99) = 30.99$, $p < .001$				
	Number of binge episodes per week in Trimester 1	9.14	4.11	.028	[0.98, 17.31]
	Infant age at EEG collection	7.47	1.89	<.001	[3.72, 11.21]
	Natural log of infant EEG age	-154.65	29.61	<.001	[-213.41, -95.89]
N2 latency	Model: $n = 106$, model $R^2 = .24$, $F(3, 102) = 10.92$, $p < .001$				
	Number of binge episodes per week in Trimester 1	17.67	13.68	.199	[-9.46, 44.79]
	Infant age at EEG collection	-5.17	0.92	<.001	[-7.00, -3.35]
	Number of retained EEG trials	0.14	0.21	.516	[-0.28, 0.56]

Note. VEP = visual-evoked potential; CI = confidence interval; LL = lower limit; UL = upper limit; EEG = electroencephalography.

Table 9 for a regression table of number of binge episodes per week in the first trimester predicting each VEP component at Visit 2.

Given previous findings from Scher et al. (1998) as well as the dysmyelinating effect of PAE and the link between ERP component latency and myelination, our primary outcome of interest was in VEP latencies, to reduce the number of total comparisons performed in the study. However, given potential interest in the effects of PAE on VEP amplitudes, we include exploratory analyses of VEP amplitudes in Tables S2–S5 in the online supplemental materials. Notably, we do not find any statistically significant effects of PAE (number of drinks per week in each trimester or number of binge episodes per week in the first trimester) on VEP amplitudes (also corrected for previous component amplitude, that is, corrected P1 amplitude = P1–N1 amplitude and corrected N2 amplitude = N2–P1

amplitude) at either timepoint that survive correction for multiple comparisons in these supplemental analyses.

Exploratory Results of T1w/T2w Ratios in Relation to VEP Latencies

Myelination in the Primary Visual Cortex (V1)

Exploratory cross-sectional analyses (n before removing outliers = 47) of whether MRI-estimates of myelin related to individual differences in VEP latencies regardless of PAE status revealed that estimates of myelination in the primary visual cortex (V1) as indexed by the T1w/T2w ratio significantly predicted uncorrected P1 latency ($b = -12.80$, $p = .040$, 95% CI [-24.98, -0.61]) such that those with decreased myelination had longer P1 latencies, when controlling

Table 9

Regression Models of Number of Binge Episodes per Week in Each Trimester Predicting VEP Component Latencies at Visit 2 (n Before Removing Outliers = 102)

Component	Variable	<i>b</i>	<i>SE</i>	<i>p</i>	95% CI [LL, UL]
N1 latency	Model: $n = 101$, model $R^2 = .11$, $F(3, 97) = 3.94$, $p = .011$				
	Number of binge episodes per week in Trimester 1	1.20	3.72	.747	[-6.17, 8.58]
	Infant age at EEG collection	-0.29	0.09	<.001	[-0.46, -0.12]
	Number of retained EEG trials	0.01	0.04	.754	[-0.07, 0.09]
P1 latency	Model: $n = 100$, model $R^2 = .02$, $F(3, 96) = 0.56$, $p = .645$				
	Number of binge episodes per week in Trimester 1	-0.92	3.04	.763	[-6.96, 5.12]
	Infant age at EEG collection	-0.06	0.07	.414	[-0.20, 0.08]
	Number of retained EEG trials	-0.02	0.03	.485	[-0.09, 0.04]
N2 latency	Model: $n = 101$, model $R^2 = .01$, $F(3, 97) = 0.47$, $p = .702$				
	Number of binge episodes per week in Trimester 1	-11.38	11.31	.317	[-33.84, 11.07]
	Infant age at EEG collection	-0.16	0.26	.553	[-0.68, 0.36]
	Number of retained EEG trials	0.05	0.13	.694	[-0.20, 0.30]

Note. VEP = visual-evoked potential; CI = confidence interval; LL = lower limit; UL = upper limit; EEG = electroencephalography.

for infant age at MRI collection, infant age at EEG collection, and number of retained VEP trials ($n = 45$, model $R^2 = .63$, $F(4, 40) = 16.82$, $p < .001$; see Figure 8 for residual plot). Note uncorrected latencies were used for these analyses as the underlying visual regions contribute to multiple VEP components' latencies simultaneously. V1 myelination indexed by the T1w/T2w ratio did not significantly predict either N1 latency (n after removing outliers = 46, model $R^2 = .34$, $F(4, 41) = 5.39$, $p = .001$) or uncorrected N2 latency ($n = 46$, model $R^2 = .27$, $F(4, 41) = 3.72$, $p = .011$). See Table 10 for a regression table of primary visual cortex (V1) T1w/T2w estimates of myelin predicting each VEP component.

Myelination in the Middle Temporal Visual Area (MT/V5)

Estimates of myelination in the middle temporal visual area (i.e., MT or V5) did not significantly predict latency at any VEP component it serves (i.e., P1 or N2; Barnikol et al., 2006) when controlling for infant age at MRI collection, infant age at EEG collection, and number of retained VEP trials (n before removing outliers = 47; P1 model: $n = 45$, model $R^2 = .61$, $F(4, 40) = 15.35$, $p < .001$; N2 model: $n = 46$, model $R^2 = .26$, $F(4, 41) = 3.55$, $p = .014$). See Table 11 for a regression table of middle temporal visual area (MT/V5) T1w/T2w estimates of myelin predicting the P1 and N2 VEP components.

Discussion


Though a rich literature has linked PAE to visually-mediated sensory and behavioral differences over childhood (Hunt et al., 1995; Jacobson et al., 1993; Janzen, 1995; Mattson et al., 1996, 1998; Uecker & Nadel, 1996), there is still a paucity of research exploring neurodevelopmental mechanisms in humans that may drive such behavioral differences (Coles & Li, 2011; Donald et al., 2015; Wozniak & Muetzel, 2011). Here we leveraged a multimodal longitudinal study to address this gap by examining early visual

neurodevelopment as a candidate pathway. We observed differential effects of PAE on visual neurodevelopment according to the following predictor characteristics: trimester of exposure, dose, and drinking pattern (i.e., presence of binge episodes). In this longitudinal sample, PAE differentially impacted the EEG-derived VEP P1 component measured within the first 6 postnatal months of age, but these effects resolved temporally by the end of the first postnatal year. In a separate analysis in the subsample of infants (collapsing across those with and without PAE) with MRI measures modeled to be sensitive to myelin in visual cortical regions known to drive the VEP response, we also established an association between individual differences in myelination and the VEP response latencies over the first postnatal year of life. These findings provide evidence that PAE and postnatal myelin may each alter early visual neurodevelopmental timing over infancy in the following ways.

First, we found that increased PAE resulted in less mature electrophysiological responses to visual stimuli, as indexed by longer VEP component latencies, depending on trimester of exposure, specific VEP component, and drinking pattern. Specifically, we found that increased alcohol exposure in the first trimester resulted in longer latency of the P1 component assessed within the first 6 months of life, replicating and extending a prior finding from Scher and colleagues that first trimester alcohol exposure resulted in longer P1 latencies at 1 month of age (1998). A similar effect of binge drinking on VEP P1 latencies was found as well, such that increased exposure to binge drinking in the first trimester resulted in increased latencies of the P1 component over the first 6 postnatal months of life. These effects together suggest that the P1 component is particularly sensitive to PAE exposure. These differential effects of trimester and consistent findings across both binge and nonbinge-patterned drinking provide further evidence that timing and drinking patterns are important in characterizing the effects of PAE.

The first trimester is a time of rapid visual development for the fetus as the building blocks of visual structures and many connections are established during this period (Finlay & Darlington,

Figure 8
Residual Plot for Primary Visual Cortex (V1) T1w/T2w Ratios (Myelin Estimates) Predicting P1 Latency ($n = 45$)



Note. T1w = T1-weighted; T2w = T2-weighted. See the online article for the color version of this figure.

Table 10

Regression Models of Primary Visual Cortex (V1) T1w/T2w Ratios (Myelin Estimates) Predicting VEP Component Latencies (n Before Removing Outliers = 47)

Component	Variable	<i>b</i>	<i>SE</i>	<i>p</i>	95% CI [LL, UL]
N1 latency	Model: $n = 46$, model $R^2 = .34$, $F(4, 41) = 5.39$, $p = .001$				
	Primary visual cortex (V1) T1w/T2w ratio	−3.59	6.29	.571	[−16.30, 9.11]
	Infant age at EEG collection	−0.25	0.24	.308	[−0.74, 0.24]
	Infant age at MRI collection	−0.25	0.26	.328	[−0.77, 0.27]
	Number of retained EEG trials	−0.12	0.08	.160	[−0.28, 0.05]
P1 latency	Model: $n = 45$, model $R^2 = .63$, $F(4, 40) = 16.82$, $p < .001$				
	Primary visual cortex (V1) T1w/T2w ratio	−12.80	6.03	.040	[−24.98, −0.61]
	Infant age at EEG collection	−0.18	0.24	.444	[−0.66, 0.30]
	Infant age at MRI collection	−0.71	0.25	.007	[−1.22, −0.20]
	Number of retained EEG trials	−0.06	0.08	.445	[−0.22, 0.10]
N2 latency	Model: $n = 46$, model $R^2 = .27$, $F(4, 41) = 3.72$, $p = .011$				
	Primary visual cortex (V1) T1w/T2w ratio	−49.48	39.56	.218	[−129.38, 30.43]
	Infant age at EEG collection	−1.58	1.51	.304	[−4.63, 1.48]
	Infant age at MRI collection	−1.06	1.61	.514	[−4.32, 2.20]
	Number of retained EEG trials	−0.23	0.52	.658	[−1.27, 0.81]

Note. VEP = visual-evoked potential; CI = confidence interval; LL = lower limit; UL = upper limit; EEG = electroencephalography; MRI = magnetic resonance imaging; T1w = T1-weighted; T2w = T2-weighted.

1995; Finlay et al., 2003; Hevner, 2000; Johnson, 2011). Indeed, the fetus's eyes open during the second trimester after much of the neural structural organization is completed in the first trimester (though there is substantial functional and structural revision postnatally as well). Therefore, this period of rapid development and change is likely highly vulnerable to environmental disturbances and teratogenic effects such as the PAE demonstrated in this sample and as noted in other samples (e.g., Scher et al., 1998).

Furthermore, the pattern of findings across the two longitudinal study visits suggest that these early VEP latency differences may manifest in developmental delays that resolve neurally by the end of the first year. Specifically, analyses of the Visit 1 EEG data over the first 6 postnatal months also revealed significant age-related changes in latency timing such that latencies shortened with older ages. This finding that P1 VEP latencies speed up with age is consistent with prior literature on the development of the VEP over the first

years of life (Lippé et al., 2006, 2009). Our findings that PAE results in longer latencies during this developmental window of change provide one indicator of delayed development. Importantly, the lack of PAE effects on latencies in the longitudinal sample at the second visit further suggests developmental delays in basic visual processing that resolve neurally by the end of the first postnatal year. Although these VEP differences were no longer observable within the longitudinal sample after 6 months of age, this should not be taken to indicate that earlier VEP delays have no developmental consequences. Altered neurodevelopmental timing, especially within sensory cortices, can have critical and lasting consequences for cognition and behavior, and mistiming of neurodevelopment is implicated in a variety of other conditions and disorders (e.g., Hensch, 2005; Ivanov, 2021; Makinodan et al., 2012; Marín, 2016; Takesian & Hensch, 2013). Moreover, visually mediated sensory and behavioral differences following PAE have been documented from childhood through

Table 11

Regression Models of Middle Temporal Visual Area (MT/V5) T1w/T2w Ratios (Myelin Estimates) Predicting Relevant VEP Component Latencies (n Before Removing Outliers = 47)

Component	Variable	<i>b</i>	<i>SE</i>	<i>p</i>	95% CI [LL, UL]
P1 latency	Model: $n = 45$, model $R^2 = .61$, $F(4, 40) = 15.35$, $p < .001$				
	Middle temporal visual area (MT/V5) T1w/T2w ratio	−10.97	7.63	.158	[−26.38, 4.44]
	Infant age at EEG collection	−0.19	0.24	.440	[−0.69, 0.30]
	Infant age at MRI collection	−0.70	0.26	.010	[−1.22, −0.18]
	Number of retained EEG trials	−0.06	0.08	.436	[−0.23, 0.10]
N2 latency	Model: $n = 46$, model $R^2 = .26$, $F(4, 41) = 3.55$, $p = .014$				
	Middle temporal visual area (MT/V5) T1w/T2w ratio	−50.14	48.76	.310	[−148.61, 48.33]
	Infant age at EEG collection	−1.57	1.53	.309	[−4.65, 1.51]
	Infant age at MRI collection	−1.02	1.62	.532	[−4.30, 2.26]
	Number of retained EEG trials	−0.25	0.52	.637	[−1.30, 0.81]

Note. VEP = visual-evoked potential; CI = confidence interval; LL = lower limit; UL = upper limit; EEG = electroencephalography; MRI = magnetic resonance imaging; T1w = T1-weighted; T2w = T2-weighted.

adulthood (Hunt et al., 1995; Jacobson et al., 1993; Janzen, 1995; Mattson et al., 1996, 1998; Uecker & Nadel, 1996), consistent with longer term consequences of early neurodevelopmental changes. Whether the currently observed pattern of delayed VEP development contributes to behavioral differences within this sample remains to be tested at later visits within this longitudinal study.

To explore one potential structural mechanism that could contribute to VEP latencies in development (independent of PAE), we also tested associations in a subsample of infants with an MRI-derived estimate of myelin, the T1w/T2w ratio. PAE has been shown repeatedly to impact myelination (Darbinian & Selzer, 2022; Donald et al., 2015; Ghazi Sherbaf et al., 2019; McLachlan et al., 2019; Newville et al., 2017, 2022; Wozniak & Muetzel, 2011). Moreover, while it is largely accepted that the significant accumulation of myelin in the brain at least partially explains the shortening of VEP latencies across infancy, as far as we are aware, this is the first empirical study to test the relation between cortical myelination and VEP latencies in infancy (though one prior study found decreased subcortical myelination, indexed by diffusion tensor imaging, was linked to longer P1 VEP latencies; Dubois et al., 2008). Here we show that decreased myelination, as indexed by T1w/T2w ratios, in the primary visual cortex (V1) was associated with longer P1 component VEP latencies over the first postnatal year. This association is consistent with those previously noted in adult and clinical samples (Heidari et al., 2019; Klistorner & Graham, 2021; Walt et al., 2015). This set of findings suggests that myelination over infancy may be partially responsible for the maturation of VEP component timing during this developmental window, but also indicates other factors contribute to the increasingly efficient and quick neural responses to basic visual stimuli. Future research with larger samples can evaluate the effects of PAE on myelination and whether that influence explains the latency changes related to PAE observed here in VEP P1 latency in infancy.

The present study design bolsters interpretation of the neurodevelopmental effects of PAE observed by addressing several key limitations in prior literature examining sequelae of PAE. Specifically, through the use of a nonclinical sample, we were able to capture a wider spectrum of possible PAE with increased ecological validity for the broader population drinking during pregnancy. We were also able to examine the effects of timing, dose, and drinking pattern (i.e., binge drinking) by measuring alcohol exposure continuously over each trimester. However, we do note that in the current sample, there were insufficient binge drinking cases to model effects of binge exposure beyond trimester 1. Additionally, we specifically selected our sample for the current study to have prenatal exposure to exclusively alcohol (i.e., no comorbid substance exposure) to better elucidate the effects of PAE specifically. Prior neurodevelopmental and behavioral research in humans has largely been subject to the confounds that come with polysubstance exposure prenatally (Coles & Li, 2011; Donald et al., 2015). Finally, we selected a comparison sample that was matched to the PAE sample on maternal age at infant birth, maternal education, maternal income, and EPDS depression scores to further reduce potential sociodemographic confounds in analyses and results.

Progress in future work should consider limitations of the current study. First, this study sample had limited endorsement of drinking in later trimesters (especially when characterizing binge drinking behaviors). The limited variability in drinking endorsement in later pregnancy meant that this study was underpowered to detect effects

of alcohol exposure in the third trimester, if any exist. While this sample structure precluded careful examination of the effects of PAE in later trimesters on visual neurodevelopment, it also reflects ecologically valid sampling of the patterns found in the community. Second, future work can include larger sample sizes of infants with both MRI and electrophysiological data to facilitate more sensitive analyses of associations between MRI-derived myelin measures and EEG-derived VEP latencies as well as explicitly test how PAE may impact these associations. The current limited sample size made it difficult to examine or detect potential PAE effects on associations between the MRI and EEG measures. The limited sample size, and thus limited statistical power to detect a wide range of effects from subtle to large, also made it difficult to make specificity claims about which visual areas' myelination patterns better explain VEP component latencies. That is, some visual regions where no statistical effects were noted in this sample may demonstrate effects in a larger and more sensitive sample. Indeed, the preliminary results from this sample demonstrating an association between MRI-derived myelin and VEP latency should motivate further research in this area. Third, the current sample, though longitudinal, included only two timepoints of data per individual over a large age range for each study visit, unlike many longitudinal designs in the first year of life. Thus, while the study was able to test several nonlinear associations between VEP measures and age at the between-participant level (i.e., cross-sectionally), the study was unable to model nonlinear age associations within-participant (i.e., longitudinally). Though this design facilitated study flexibility for families and retention over the first year in this community, future studies may prioritize more visits within the first year to facilitate additional analysis strategies like nonlinear longitudinal growth models. Next, future work can more completely measure comorbid nonalcohol substance exposure over the entire period of pregnancy as opposed to the 3 months prior to assessment in pregnancy. The current strategy may have missed comorbid substance use occurring outside of the assessment window. As a result, we are not able to statistically exclude potential effects of comorbid substance exposures in early pregnancy, outside of the assessment window. That is, there may be effects of these exposures in the results presented that could not be explicitly attributed to comorbid substance use outside of the assessment window. The current sampling strategy was employed to balance participant burden as this analysis was not part of the primary aims for the ongoing longitudinal study. Finally, although the study is ongoing and in the future can examine whether behavioral consequences emerge from these early infant neurodevelopmental changes, the current set of analyses cannot link the neurodevelopmental measures to any behavioral phenotypes at this time.

Finally, with an eye toward supporting future pregnant people and infants in contexts of prenatal alcohol use and exposure, we empirically probed whether a potential screening tool would relate to self-reported alcohol use in our sample. Further supporting the robustness of the WHO ASSIST, a lifetime risk measure, we found that the alcohol risk score calculated using this tool was related to the self-report of number of drinks per week and of binge episodes during pregnancy throughout all three trimesters for this South African community. Moreover, 100% of the mothers who scored above lower risk on the WHO ASSIST reported drinking during pregnancy, so this tool may be useful to identify individuals who may benefit from counseling and support around drinking during pregnancy. This analysis is one of the first empirical evaluations of the WHO ASSIST as a tool

to detect those at higher risk of endorsing alcohol use during pregnancy and further demonstrates its utility and acceptability in an underresourced community like the Gugulethu community (WHO ASSIST Working Group, 2002).

In conclusion, prenatal exposure to alcohol, specifically in the first trimester, may alter timing of visual neurodevelopment in the first 6 months of life, with possible consequences lasting across the lifespan. Future directions of this work can identify pre- and postnatal resilience factors that may protect against the effects of PAE described in this study. Given our exploratory findings that myelin is also separately related to VEP latency in infancy (across those with and without PAE), future studies should examine whether there are links between PAE, myelination, and VEP latency that could explain the VEP differences we observed here and offer a postnatal intervention target. In these ways, better elucidating the neurodevelopmental sequelae of PAE using longitudinal multimodal studies such as the current study will be important in identifying how to best support pregnant people and their infants' development.

References

- Adebisi, B. O., Mukumbang, F. C., & Beytell, A.-M. (2021). Policy requirements for the prevention and management of fetal alcohol spectrum disorder in South Africa: A policy brief. *Frontiers in Public Health*, 9, Article 592726. <https://doi.org/10.3389/fpubh.2021.592726>
- Almeida, L., Andreu-Fernández, V., Navarro-Tapia, E., Aras-López, R., Serra-Delgado, M., Martínez, L., García-Algar, O., & Gómez-Roig, M. D. (2020). Murine models for the study of fetal alcohol spectrum disorders: An overview. *Frontiers in Pediatrics*, 8, Article 359. <https://doi.org/10.3389/fped.2020.00359>
- Amso, D., & Scerif, G. (2015). The attentive brain: Insights from developmental cognitive neuroscience. *Nature Reviews Neuroscience*, 16(10), 606–619. <https://doi.org/10.1038/nrn4025>
- Antonov, N. K., Ruzal-Shapiro, C. B., Morel, K. D., Millar, W. S., Kashyap, S., Lauren, C. T., & Garzon, M. C. (2017). Feed and wrap MRI technique in infants. *Clinical Pediatrics*, 56(12), 1095–1103. <https://doi.org/10.1177/0009922816677806>
- Bailey, B. N., Delaney-Black, V., Covington, C. Y., Ager, J., Janisse, J., Hannigan, J. H., & Sokol, R. J. (2004). Prenatal exposure to binge drinking and cognitive and behavioral outcomes at age 7 years. *American Journal of Obstetrics and Gynecology*, 191(3), 1037–1043. <https://doi.org/10.1016/j.ajog.2004.05.048>
- Barnikol, U. B., Amunts, K., Dammers, J., Mohlberg, H., Fieseler, T., Malikovic, A., Zilles, K., Niedeggen, M., & Tass, P. A. (2006). Pattern reversal visual evoked responses of V1/V2 and V5/MT as revealed by MEG combined with probabilistic cytoarchitectonic maps. *NeuroImage*, 31(1), 86–108. <https://doi.org/10.1016/j.neuroimage.2005.11.045>
- Bary-Anwar, R., Riggins, T., & Scott, L. S. (2020). Electrophysiology in developmental populations: Key methods and findings. In K. Cohen Kadosh (Ed.), *The Oxford handbook of developmental cognitive neuroscience*. Oxford University Press. <https://doi.org/10.1093/oxfordhb/9780198827474.013.3>
- CDC. (2022, November 28). *Binge drinking*. <https://www.cdc.gov/alcohol/fact-sheets/binge-drinking.htm>
- Coles, C. (1994). Critical periods for prenatal alcohol exposure. *Alcohol Health and Research World*, 18(1), 22–29.
- Coles, C. D., & Li, Z. (2011). Functional neuroimaging in the examination of effects of prenatal alcohol exposure. *Neuropsychology Review*, 21(2), 119–132. <https://doi.org/10.1007/s11065-011-9165-y>
- Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression. *British Journal of Psychiatry*, 150(6), 782–786. <https://doi.org/10.1192/bjp.150.6.782>
- Crenshaw, K. (1989). Demarginalizing the intersection of race and sex: A Black feminist critique of antidiscrimination doctrine, feminist theory and antiracist politics. *University of Chicago Legal Forum*, 1989(1), Article 8. <https://chicagounbound.uchicago.edu/uclf/vol1989/iss1/8>
- Darbinian, N., & Selzer, M. E. (2022). Oligodendrocyte pathology in fetal alcohol spectrum disorders. *Neural Regeneration Research*, 17(3), 497–502. <https://doi.org/10.4103/1673-5374.314294>
- Dean, D. C., O'Muircheartaigh, J., Dirks, H., Waskiewicz, N., Lehman, K., Walker, L., Han, M., & Deoni, S. C. L. (2014). Modeling healthy male white matter and myelin development: 3 through 60 months of age. *NeuroImage*, 84, 742–752. <https://doi.org/10.1016/j.neuroimage.2013.09.058>
- Dejong, K., Olyaei, A., & Lo, J. O. (2019). Alcohol use in pregnancy. *Clinical Obstetrics & Gynecology*, 62(1), 142–155. <https://doi.org/10.1097/GRF.0000000000000414>
- Deoni, S. C. L., Mercure, E., Blasi, A., Gasston, D., Thomson, A., Johnson, M., Williams, S. C. R., & Murphy, D. G. M. (2011). Mapping infant brain myelination with magnetic resonance imaging. *The Journal of Neuroscience*, 31(2), 784–791. <https://doi.org/10.1523/JNEUROSCI.2106-10.2011>
- Donald, K. A., Eastman, E., Howells, F. M., Adnams, C., Riley, E. P., Woods, R. P., Narr, K. L., & Stein, D. J. (2015). Neuroimaging effects of prenatal alcohol exposure on the developing human brain: A magnetic resonance imaging review. *Acta Neuropsychiatrica*, 27(5), 251–269. <https://doi.org/10.1017/neu.2015.12>
- Donald, K. A., Wedderburn, C. J., Barnett, W., Nhapi, R. T., Rehman, A. M., Stadler, J. A. M., Hoffman, N., Koen, N., Zar, H. J., & Stein, D. J. (2019). Risk and protective factors for child development: An observational South African birth cohort. *PLoS Medicine*, 16(9), Article e1002920. <https://doi.org/10.1371/journal.pmed.1002920>
- Dubois, J., Dehaene-Lambertz, G., Soarès, C., Cointepas, Y., Bihan, D. L., & Hertz-Pannier, L. (2008). Microstructural correlates of infant functional development: Example of the visual pathways. *The Journal of Neuroscience*, 28(8), 1943–1948. <https://doi.org/10.1523/JNEUROSCI.5145-07.2008>
- Finlay, B. L., Clancy, B., & Kingsbury, M. A. (2003). The developmental neurobiology of early vision. In B. Hopkins & S. P. Johnson (Eds.), *Neurobiology of infant vision* (pp. 1–41). Praeger Publishers.
- Finlay, B. L., & Darlington, R. B. (1995). Linked regularities in the development and evolution of mammalian brains. *Science*, 268(5217), 1578–1584. <https://doi.org/10.1126/science.7777856>
- Fletcher, O. V., May, P. A., Seedat, S., Sikkema, K. J., & Watt, M. H. (2018). Attitudes toward alcohol use during pregnancy among women recruited from alcohol-serving venues in Cape Town, South Africa: A mixed-methods study. *Social Science & Medicine*, 215, 98–106. <https://doi.org/10.1016/j.socscimed.2018.09.008>
- Gabard-Durnam, L., & McLaughlin, K. A. (2020). Sensitive periods in human development: Charting a course for the future. *Current Opinion in Behavioral Sciences*, 36, 120–128. <https://doi.org/10.1016/j.cobeha.2020.09.003>
- Ganzetti, M., Liu, Q., & Mantini, D. (2018). A spatial registration toolbox for structural MR imaging of the aging brain. *Neuroinformatics*, 16(2), 167–179. <https://doi.org/10.1007/s12021-018-9355-3>
- Ganzetti, M., Wenderoth, N., & Mantini, D. (2014). Whole brain myelin mapping using T1- and T2-weighted MR imaging data. *Frontiers in Human Neuroscience*, 8, Article 671. <https://doi.org/10.3389/fnhum.2014.00671>
- Ganzetti, M., Wenderoth, N., & Mantini, D. (2015). Mapping pathological changes in brain structure by combining T1- and T2-weighted MR imaging data. *Neuroradiology*, 57(9), 917–928. <https://doi.org/10.1007/s00234-015-1550-4>
- Ghazi Sherbaf, F., Aarabi, M. H., Hosein Yazdi, M., & Haghshomar, M. (2019). White matter microstructure in fetal alcohol spectrum disorders: A systematic review of diffusion tensor imaging studies. *Human Brain Mapping*, 40(3), 1017–1036. <https://doi.org/10.1002/hbm.24409>
- Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., & Ugurbil, K. (2016). A multi-modal parcellation of human

- cerebral cortex. *Nature*, 536(7615), 171–178. <https://doi.org/10.1038/nature18933>
- Glasser, M. F., & Van Essen, D. C. (2011). Mapping human cortical areas in vivo based on myelin content as revealed by T1- and T2-weighted MRI. *The Journal of Neuroscience*, 31(32), 11597–11616. <https://doi.org/10.1523/JNEUROSCI.2180-11.2011>
- Goddings, A. L., Roalf, D., Lebel, C., & Tamnes, C. K. (2021). Development of white matter microstructure and executive functions during childhood and adolescence: A review of diffusion MRI studies. *Developmental Cognitive Neuroscience*, 51, Article 101008. <https://doi.org/10.1016/j.dcn.2021.101008>
- Harter, M. R., Deaton, F. K., & Odom, J. V. (1977). Maturation of evoked potentials and visual preference in 6–45-day-old infants: Effects of check size, visual acuity, and refractive error. *Electroencephalography and Clinical Neurophysiology*, 42(5), 595–607. [https://doi.org/10.1016/0013-4694\(77\)90278-4](https://doi.org/10.1016/0013-4694(77)90278-4)
- Hartmann, E. E. (1995). Infant visual development: An overview of studies using visual evoked potential measures from Harter to the present. *International Journal of Neuroscience*, 80(1–4), 203–235. <https://doi.org/10.3109/00207459508986101>
- Heidari, M., Radcliff, A. B., McLellan, G. J., Ver Hoeve, J. N., Chan, K., Kiland, J. A., Keuler, N. S., August, B. K., Sebo, D., Field, A. S., & Duncan, I. D. (2019). Evoked potentials as a biomarker of remyelination. *Proceedings of the National Academy of Sciences*, 116(52), 27074–27083. <https://doi.org/10.1073/pnas.1906358116>
- Hensch, T. K. (2005). Critical period plasticity in local cortical circuits. *Nature Reviews Neuroscience*, 6(11), 877–888. <https://doi.org/10.1038/nrn1787>
- Hensch, T. K., & Bilimoria, P. M. (2012). Re-opening windows: Manipulating critical periods for brain development. *Cerebrum: The Dana Forum on Brain Science*, 2012, Article 11.
- Hevner, R. F. (2000). Development of connections in the human visual system during fetal mid-gestation: A DiI-tracing study. *Journal of Neuropathology & Experimental Neurology*, 59(5), 385–392. <https://doi.org/10.1093/jnen/59.5.385>
- Ho, D. E., Imai, K., King, G., & Stuart, E. A. (2011). Matchit: Nonparametric preprocessing for parametric causal inference. *Journal of Statistical Software*, 42(8), 1–28. <https://doi.org/10.18637/jss.v042.i08>
- Hunt, E., Streissguth, A. P., Kerr, B., & Olson, H. C. (1995). Mothers' alcohol consumption during pregnancy: Effects on spatial-visual reasoning in 14-year-old children. *Psychological Science*, 6(6), 339–342. <https://doi.org/10.1111/j.1467-9280.1995.tb00523.x>
- Ivanov, P. C. (2021). The new field of network physiology: Building the human physiome. *Frontiers in Network Physiology*, 1, Article 711778. <https://doi.org/10.3389/fnetp.2021.711778>
- Jacobson, J. L., Akkaya-Hocagil, T., Ryan, L. M., Dodge, N. C., Richardson, G. A., Olson, H. C., Coles, C. D., Day, N. L., Cook, R. J., & Jacobson, S. W. (2021). Effects of prenatal alcohol exposure on cognitive and behavioral development: Findings from a hierarchical meta-analysis of data from six prospective longitudinal U. S. cohorts. *Alcoholism: Clinical and Experimental Research*, 45(10), 2040–2058. <https://doi.org/10.1111/acer.14686>
- Jacobson, S. W., Jacobson, J. L., Molteno, C. D., Warton, C. M. R., Wintermark, P., Hoyme, H. E., De Jong, G., Taylor, P., Warton, F., Lindinger, N. M., Carter, R. C., Dodge, N. C., Grant, E., Warfield, S. K., Zöllei, L., van der Kouwe, A. J. W., & Meintjes, E. M. (2017). Heavy prenatal alcohol exposure is related to smaller corpus callosum in newborn MRI scans. *Alcoholism: Clinical and Experimental Research*, 41(5), 965–975. <https://doi.org/10.1111/acer.13363>
- Jacobson, S. W., Jacobson, J. L., Sokol, R. J., Martier, S. S., & Ager, J. W. (1993). Prenatal alcohol exposure and infant information processing ability. *Child Development*, 64(6), 1706–1721. <https://doi.org/10.2307/1131464>
- Janzen, L. (1995). Neuropsychological evaluation of preschoolers with fetal alcohol syndrome. *Neurotoxicology and Teratology*, 17(3), 273–279. [https://doi.org/10.1016/0892-0362\(94\)00063-J](https://doi.org/10.1016/0892-0362(94)00063-J)
- Johnson, S. P. (2011). Development of visual perception. *Wiley Interdisciplinary Reviews: Cognitive Science*, 2(5), 515–528. <https://doi.org/10.1002/wcs.128>
- Jones, K. L., & Smith, D. W. (1973). Recognition of the fetal alcohol syndrome in early infancy. *The Lancet*, 302(7836), 999–1001. [https://doi.org/10.1016/S0140-6736\(73\)91092-1](https://doi.org/10.1016/S0140-6736(73)91092-1)
- Kelly, S. J., Pierce, D. R., & West, J. R. (1987). Microencephaly and hyperactivity in adult rats can be induced by neonatal exposure to high blood alcohol concentrations. *Experimental Neurology*, 96(3), 580–593. [https://doi.org/10.1016/0014-4886\(87\)90220-2](https://doi.org/10.1016/0014-4886(87)90220-2)
- Klistorner, A., & Graham, S. L. (2021). Role of multifocal visually evoked potential as a biomarker of demyelination, spontaneous remyelination, and myelin repair in multiple sclerosis. *Frontiers in Neuroscience*, 15, Article 725187. <https://doi.org/10.3389/fnins.2021.725187>
- Kwiatkowski, M. A., Donald, K. A., Stein, D. J., Ipser, J., Thomas, K. G. F., & Roos, A. (2018). Cognitive outcomes in prenatal methamphetamine exposed children aged six to seven years. *Comprehensive Psychiatry*, 80, 24–33. <https://doi.org/10.1016/j.comppsych.2017.08.003>
- Lawrie, T. A., Hofmeyr, G. J., De Jager, M., & Berk, M. (1998). Validation of the Edinburgh Postnatal Depression Scale on a cohort of South African women. *South African Medical Journal*, 88(10), 1340–1344.
- Lees, B., Mewton, L., Jacobus, J., Valadez, E. A., Stapinski, L. A., Teesson, M., Tapert, S. F., & Squeglia, L. M. (2020). Association of prenatal alcohol exposure with psychological, behavioral, and neurodevelopmental outcomes in children from the adolescent brain cognitive development study. *American Journal of Psychiatry*, 177(11), 1060–1072. <https://doi.org/10.1176/appi.ajp.2020.20010086>
- Lemoine, P., Harousseau, H., Borleyru, J. P., & Menuet, J. C. (1968). Les enfants de parents alcooliques: Anomalies, observées de 127 cas [Children of alcoholic parents: Anomalies observed in 127 cases]. *Quest Medical*, 21, 476–482.
- Lippé, S., Kovacevic, N., & McIntosh, R. (2009). Differential maturation of brain signal complexity in the human auditory and visual system. *Frontiers in Human Neuroscience*, 3, Article 48. <https://doi.org/10.3389/neuro.09.048.2009>
- Lippé, S., Roy, M.-S., Perchet, C., & Lassonde, M. (2006). Electrophysiological markers of visuocortical development. *Cerebral Cortex*, 17(1), 100–107. <https://doi.org/10.1093/cercor/bhj130>
- Makinodan, M., Rosen, K. M., Ito, S., & Corfas, G. (2012). A critical period for social experience-dependent oligodendrocyte maturation and myelination. *Science*, 337(6100), 1357–1360. <https://doi.org/10.1126/science.1220845>
- Margolis, E. (2024, January 15). *Longitudinal effects of prenatal alcohol exposure on visual neurodevelopment over infancy*. https://osf.io/kjtsu/?view_only=30b08cd27a0f412cbff10dddf39561fd
- Marin, O. (2016). Developmental timing and critical windows for the treatment of psychiatric disorders. *Nature Medicine*, 22(11), 1229–1238. <https://doi.org/10.1038/nm.4225>
- Marshall, A. T., Bodison, S. C., Uban, K. A., Adise, S., Jonker, D., Charles, W., Donald, K. A., Kan, E., Ipser, J. C., Butler-Kruger, L., Steigelmann, B., Narr, K. L., Joshi, S. H., Brink, L. T., Odendaal, H. J., Scheffler, F., Stein, D. J., & Sowell, E. R. (2022). The impact of prenatal alcohol and/or tobacco exposure on brain structure in a large sample of children from a South African birth cohort. *Alcoholism: Clinical and Experimental Research*, 46(11), 1980–1992. <https://doi.org/10.1111/acer.14945>
- Mattson, S. N., Gramling, L., Riley, E. P., Delis, D. C., & Jones, K. L. (1996). Global—local processing in children prenatally exposed to alcohol. *Child Neuropsychology*, 2(3), 165–175. <https://doi.org/10.1080/09297049608402249>
- Mattson, S. N., Riley, E. P., Gramling, L., & Delis, D. C. (1998). Neuropsychological comparison of alcohol-exposed children with or without physical features of fetal alcohol syndrome. *Neuropsychology*, 12(1), 146–153. <https://doi.org/10.1037/0894-4105.12.1.146>

- May, P. A., Marais, A.-S., De Vries, M., Hasken, J. M., Stegall, J. M., Hedrick, D. M., Snell, C. L., Seedat, S., & Parry, C. D. H. (2019). The dop system of alcohol distribution is dead, but it's legacy lives on.... *International Journal of Environmental Research and Public Health*, 16(19), Article 3701. <https://doi.org/10.3390/ijerph16193701>
- Mazerolle, M. J. (2023). *AICcmodavg: Model selection and multimodel inference based on (Q)AIC(c)*. <https://cran.r-project.org/package=AICcmodavg>
- McCulloch, D. L. (2013). Visual evoked potentials in infants. In M. De Haan (Ed.), *Infant EEG and event-related potentials* (pp. 39–76). Psychology Press.
- McLachlan, K., Vavasour, I., MacKay, A., Brain, U., Oberlander, T., Loock, C., Reynolds, J. N., & Beaulieu, C. (2019). Myelin water fraction imaging of the brain in children with prenatal alcohol exposure. *Alcoholism: Clinical and Experimental Research*, 43(5), 833–841. <https://doi.org/10.1111/acer.14024>
- Milbocker, K. A., LeBlanc, G. L., Brengel, E. K., Hekmatyar, K. S., Kulkarni, P., Ferris, C. F., & Klintoova, A. Y. (2022). Reduced and delayed myelination and volume of corpus callosum in an animal model of Fetal Alcohol Spectrum Disorders partially benefit from voluntary exercise. *Scientific Reports*, 12(1), Article 10653. <https://doi.org/10.1038/s41598-022-14752-3>
- Monachino, A. D., Lopez, K. L., Pierce, L. J., & Gabard-Durnam, L. J. (2022). The HAPPE plus Event-Related (HAPPE + ER) software: A standardized preprocessing pipeline for event-related potential analyses. *Developmental Cognitive Neuroscience*, 57, Article 101140. <https://doi.org/10.1016/j.dcn.2022.101140>
- National Institute on Alcohol Abuse and Alcoholism. (2000). *10th Special Report to the U.S. Congress on Alcohol and Health*. <https://pubs.niaaa.nih.gov/publications/10report/10thspecialreport.pdf>
- Nelson, C. A., & Gabard-Durnam, L. J. (2020). Early adversity and critical periods: Neurodevelopmental consequences of violating the expectable environment. *Trends in Neurosciences*, 43(3), 133–143. <https://doi.org/10.1016/j.tins.2020.01.002>
- Newville, J., Howard, T. A., Chavez, G. J., Valenzuela, C. F., & Cunningham, L. A. (2022). Persistent myelin abnormalities in a third trimester-equivalent mouse model of fetal alcohol spectrum disorder. *Alcoholism: Clinical and Experimental Research*, 46(1), 77–86. <https://doi.org/10.1111/acer.14752>
- Newville, J., Valenzuela, C. F., Li, L., Jantzie, L. L., & Cunningham, L. A. (2017). Acute oligodendrocyte loss with persistent white matter injury in a third trimester equivalent mouse model of fetal alcohol spectrum disorder. *Glia*, 65(8), 1317–1332. <https://doi.org/10.1002/glia.23164>
- Nulman, I., Rovet, J., Kennedy, D., Wasson, C., Gladstone, J., Fried, S., & Koren, G. (2004). Binge alcohol consumption by non-alcohol—dependent women during pregnancy affects child behaviour, but not general intellectual functioning: A prospective controlled study. *Archives of Women's Mental Health*, 7(3), 173–181. <https://doi.org/10.1007/s00737-004-0055-7>
- O'Leary, C. M., Nassar, N., Zubrick, S. R., Kurinczuk, J. J., Stanley, F., & Bower, C. (2010). Evidence of a complex association between dose, pattern and timing of prenatal alcohol exposure and child behaviour problems. *Addiction*, 105(1), 74–86. <https://doi.org/10.1111/j.1360-0443.2009.02756.x>
- Olegård, R., Sabel, K.-G., Aronsson, M., Sandin, B., Johansson, P. R., Carlsson, C., Kyllerman, M., Iversen, K., & Hrbek, A. (1979). Effects on the child of alcohol abuse during pregnancy: Retrospective and prospective studies. *Acta Paediatrica*, 68(s275), 112–121. <https://doi.org/10.1111/j.1651-2227.1979.tb06170.x>
- Pellowski, J. A., Wedderburn, C. J., Groenewold, N. A., Roos, A., Subramoney, S., Hoffman, N., Fouché, J.-P., Joshi, S. H., Woods, R. P., Narr, K. L., Zar, H. J., Donald, K. A., & Stein, D. J. (2023). Maternal perinatal depression and child brain structure at 2-3 years in a South African birth cohort study. *Translational Psychiatry*, 13(1), Article 96. <https://doi.org/10.1038/s41398-023-02395-5>
- Petersen Williams, P., Jordaan, E., Mathews, C., Lombard, C., & Parry, C. D. H. (2014). Alcohol and other drug use during pregnancy among women attending midwife obstetric units in the Cape Metropole, South Africa. *Advances in Preventive Medicine*, 2014, Article 871427. <https://doi.org/10.1155/2014/871427>
- Pini, N., Shuffrey, L. C., Lucchini, M., Sania, A., Nelson, M. E., David Nugent, J., Ochoa, T. N., Odendaal, H. J., Fifer, W. P., Myers, M. M., & Elliott, A. J. (2019, July). *Cluster analysis of alcohol consumption during pregnancy in the safe passage study*. [Conference session] In 2019 41st Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC) (pp. 1338–1341). IEEE.
- Popova, S., Charness, M. E., Burd, L., Crawford, A., Hoyme, H. E., Mukherjee, R. A. S., Riley, E. P., & Elliott, E. J. (2023). Fetal alcohol spectrum disorders. *Nature Reviews Disease Primers*, 9(1), Article 11. <https://doi.org/10.1038/s41572-023-00420-x>
- Popova, S., Lange, S., Probst, C., Gmel, G., & Rehm, J. (2018). Global prevalence of alcohol use and binge drinking during pregnancy, and fetal alcohol spectrum disorder. *Biochemistry and Cell Biology*, 96(2), 237–240. <https://doi.org/10.1139/bcb-2017-0077>
- Pretorius, L., Naidoo, A., & Reddy, S. P. (2009). “Kitchen Cupboard Drinking”: A review of South African women's secretive alcohol addiction, treatment history, and barriers to accessing treatment. *Social Work in Public Health*, 24(1–2), 89–99. <https://doi.org/10.1080/19371910802569575>
- R Core Team. (2022). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Reh, R. K., Dias, B. G., Nelson, C. A., Kaufer, D., Werker, J. F., Kolb, B., Levine, J. D., & Hensch, T. K. (2020). Critical period regulation across multiple timescales. *Proceedings of the National Academy of Sciences*, 117(38), 23242–23251. <https://doi.org/10.1073/pnas.1820836117>
- Rosen, M. L., Amso, D., & McLaughlin, K. A. (2019). The role of the visual association cortex in scaffolding prefrontal cortex development: A novel mechanism linking socioeconomic status and executive function. *Developmental Cognitive Neuroscience*, 39, Article 100699. <https://doi.org/10.1016/j.dcn.2019.100699>
- RStudio Team. (2015). *RStudio: Integrated development environment for R*. <https://www.rstudio.com/>
- Scher, M. S., Richardson, G. A., Robles, N., Geva, D., Goldschmidt, L., Dahl, R. E., Scabassi, R. J., & Day, N. L. (1998). Effects of prenatal substance exposure: Altered maturation of Visual Evoked Potentials. *Pediatric Neurology*, 18(3), 236–243. [https://doi.org/10.1016/S0887-8994\(97\)00217-8](https://doi.org/10.1016/S0887-8994(97)00217-8)
- Signorelli, A. (2023). *DescTools: Tools for descriptive statistics*. <https://CRAN.R-project.org/package=DescTools>
- Smith, I. E., Coles, C. D., Lancaster, J., Fernhoff, P. M., & Falek, A. (1986). The effect of volume and duration of prenatal ethanol exposure on neonatal physical and behavioral development. *Neurobehavioral Toxicology and Teratology*, 8(4), 375–381.
- Stein, D. J., Koen, N., Donald, K. A., Adnams, C. M., Koopowitz, S., Lund, C., Marais, A., Myers, B., Roos, A., Sorsdahl, K., Stern, M., Tomlinson, M., van der Westhuizen, C., Vythilingum, B., Myer, L., Barnett, W., Brittain, K., & Zar, H. J. (2015). Investigating the psychosocial determinants of child health in Africa: The Drakenstein Child Health Study. *Journal of Neuroscience Methods*, 252, 27–35. <https://doi.org/10.1016/j.jneumeth.2015.03.016>
- Streissguth, A. P., Sampson, P. D., & Barr, H. M. (1989). Neurobehavioral dose-response effects of prenatal alcohol exposure in humans from infancy to adulthood. *Annals of the New York Academy of Sciences*, 562(1), 145–158. <https://doi.org/10.1111/j.1749-6632.1989.tb21013.x>
- Subramoney, S., Eastman, E., Adnams, C., Stein, D. J., & Donald, K. A. (2018). The early developmental outcomes of prenatal alcohol exposure: A review. *Frontiers in Neurology*, 9, Article 1108. <https://doi.org/10.3389/fneur.2018.01108>
- Takesian, A. E., & Hensch, T. K. (2013). Balancing plasticity/stability across brain development. In M. M. Merzenich, M. Nahum, & T. M. Van Vleet (Eds.), *Progress in brain research* (Vol. 207, pp. 3–34). Elsevier. <https://doi.org/10.1016/B978-0-444-63327-9.00001-1>

- Taylor, M. J., Murphy, W. J., & Whyte, H. E. (1992). Prognostic reliability of somatosensory and visual evoked potentials of asphyxiated term infants. *Developmental Medicine & Child Neurology*, 34(6), 507–515. <https://doi.org/10.1111/j.1469-8749.1992.tb11471.x>
- Testa, M., Quigley, B. M., & Eiden, R. D. (2003). The effects of prenatal alcohol exposure on infant mental development: A meta-analytical review. *Alcohol and Alcoholism*, 38(4), 295–304. <https://doi.org/10.1093/alcalc/agg087>
- The Association for Alcohol Responsibility and Education. (n.d.). *A unit of alcohol*. <https://aware.org.za/a-unit-of-alcohol/>
- Tsai, A. C., Scott, J. A., Hung, K. J., Zhu, J. Q., Matthews, L. T., Psaros, C., & Tomlinson, M. (2013). Reliability and validity of instruments for assessing perinatal depression in African settings: Systematic review and meta-analysis. *PLoS ONE*, 8(12), Article e82521. <https://doi.org/10.1371/journal.pone.0082521>
- Uecker, A., & Nadel, L. (1996). Spatial locations gone awry: Object and spatial memory deficits in children with fetal alcohol syndrome. *Neuropsychologia*, 34(3), 209–223. [https://doi.org/10.1016/0028-3932\(95\)00096-8](https://doi.org/10.1016/0028-3932(95)00096-8)
- Venables, W. N., & Ripley, B. D. (2003). *Modern applied statistics with S* (4th ed.). Springer.
- Walt, A., van der Kolbe, S., Mitchell, P., Wang, Y., Butzkueven, H., Egan, G., Yiannikas, C., Graham, S., Kilpatrick, T., & Klistorner, A. (2015). Parallel changes in structural and functional measures of optic nerve myelination after optic neuritis. *PLoS ONE*, 10(5), Article e0121084. <https://doi.org/10.1371/journal.pone.0121084>
- Watt, M. H., Eaton, L. A., Choi, K. W., Velloza, J., Kalichman, S. C., Skinner, D., & Sikkema, K. J. (2014). “It’s better for me to drink, at least the stress is going away”: Perspectives on alcohol use during pregnancy among South African women attending drinking establishments. *Social Science & Medicine*, 116, 119–125. <https://doi.org/10.1016/j.socscimed.2014.06.048>
- Werchan, D. M., & Amso, D. (2017). A novel ecological account of prefrontal cortex functional development. *Psychological Review*, 124(6), 720–739. <https://doi.org/10.1037/rev0000078>
- West, J. R., Kelly, S. J., & Pierce, D. R. (1987). Severity of alcohol-induced deficits in rats during the third trimester equivalent is determined by the pattern of exposure. *Alcohol and Alcoholism Supplement*, 1, 461–465.
- WHO ASSIST Working Group. (2002). The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): Development, reliability and feasibility. *Addiction*, 97(9), 1183–1194. <https://doi.org/10.1046/j.1360-0443.2002.00185.x>
- Williams, D. R., & Wyatt, R. (2015). Racial bias in health care and health: Challenges and opportunities. *JAMA*, 314(6), 555–556. <https://doi.org/10.1001/jama.2015.9260>
- Wozniak, J. R., & Muetzel, R. L. (2011). What does diffusion tensor imaging reveal about the brain and cognition in fetal alcohol spectrum disorders? *Neuropsychology Review*, 21(2), 133–147. <https://doi.org/10.1007/s11065-011-9162-1>
- Zieff, M. R., Miles, M., Eastman, E., Ginnell, L., Williams, S. C. R., Jones, D. K., Alexander, D. C., Gabard-Durnam, L. J., Bosco, C., McCormick, S., Klepac-Ceraj, V., Bonham, K. S., McCann, S., Patel, F., Mbale, E., Gladstone, M., Pini, N., Lucchini, M., Deoni, S. ... Donald, K. A. (under review). *Characterising developing executive functions in the first 1000 days: The Khula Cohort*.

Received June 16, 2023

Revision received January 10, 2024

Accepted January 10, 2024 ■